

Review

Dietary Polyphenols and Their Biological Significance

Xiuzhen Han, Tao Shen and Hongxiang Lou*

Department of Natural Product Chemistry, School of Pharmaceutical Sciences, Shandong University, 44 West Wenhua Road, Jinan 250012, P. R. China; E-mails: xzyhan@sina.com (X. Han); v.shentao@yahoo.com.cn (T. Shen)

* Author to whom correspondence should be addressed; E-mail: louhongxiang@sdu.edu.cn; Tel.: (+86)-531-88382012; Fax: (+86)-531-88382019

Received: 12 June 2007; in revised form: 27 August 2007 / Accepted: 27 August 2007 /

Published: 12 September 2007

1. Introduction

Oxidative stress results in oxidative alteration of biological macromolecules such as lipids, proteins and nucleic acids. It is considered to play a pivotal role in the pathogenesis of aging and degenerative diseases [1-3]. In order to cope with an excess of free radicals produced upon oxidative stress, human bodies have developed sophisticated mechanisms for maintaining redox homeostasis. These protective mechanisms include scavenging or detoxification of reactive oxygen species (ROS), blocking ROS production, sequestration of transition metals, as well as enzymatic and nonenzymatic antioxidant defenses produced in the body, that is, endogenous [4,5], and others supplied with the diet, namely, exogenous ones. Among them, dietary polyphenols have been widely studied for their strong antioxidant capacities and other properties by which cell functions are regulated [6,7].

Dietary polyphenols represent a group of secondary metabolites which widely occur in fruits, vegetables, wine, tea, extra virgin olive oil, chocolate and other cocoa products. They are mostly derivatives, and/or isomers of flavones, isoflavones, flavonols, catechins, and phenolic acids. Dietary polyphenols exhibit many biologically significant functions, such as protection against oxidative stress, and degenerative diseases. Experimental data indicate that most of these biological actions can be attributed to their intrinsic antioxidant capabilities. Dietary polyphenols may offer an indirect protection by activating endogenous defense systems and by modulating cellular signaling processes such as NF- κ B activation, AP-1 DNA binding, glutathione biosynthesis, PI3-kinase/Akt pathway, MAPK proteins (ERK, JNK and P38) activation, and the translocation into the nucleus of Nrf2 [8-10].

2. Classification and occurrence of dietary polyphenols

Dietary polyphenols are the most abundant antioxidants in human diets. With over 8,000 structural variants, they are secondary metabolites of plants and denote many substances with aromatic ring(s) bearing one or more hydroxyl moieties. They are subdivided into groups (Figure 1) by the number of phenolic rings and of the structural elements that link these rings [11]: (1) The phenolic acids with the subclasses derived from hydroxybenzoic acids such as gallic acid and from hydroxycinnamic acid, containing caffeic, ferulic, and coumaric acid; (2) the large flavonoid subclass, which includes the flavonols, flavones, isoflavones, flavanones, anthocyanidins, and flavanols; (3) the stilbenes; and (4) the lignans and the polymeric lignins.


The main dietary sources of polyphenols include some common fruits, vegetables and beverages. Phenolic acids account for about one third of the total intake and flavonoids account for the remaining two thirds. The most abundant flavonoids in the diet are flavanols (catechins plus proanthocyanidins), anthocyanins and their oxidation products. The main polyphenol dietary sources are fruit and beverages (fruit juice, wine, tea, coffee, chocolate and beer) and, to a lesser extent vegetables, dry legumes and cereals. Most of dietary polyphenols and their sources in our diets were shown in Table 1.

2.1 Phenolic acids

A major class within the phenolic compounds is the hydroxycinnamic acids, which are widely distributed in plant kingdom. The major hydroxycinnamic acid is caffeic acid, which occurs in foods mainly as an ester with quinic acid called chlorogenic acid (5-caffeoylquinic acid). Chlorogenic acid

and caffeic acid are antioxidants *in vitro* and they might inhibit the formation of mutagenic and carcinogenic *N*-nitroso compounds for the inhibitory effect on the *N*-nitrosation reaction *in vitro*.

Figure 1. Classification of dietary polyphenols.



2.2 Flavonoids

Flavonoids are the most abundant polyphenols in human diets, and are mainly divided into: (a) anthocyanins, glycosylated derivative of anthocyanidin, present in colorful flowers and fruits; (b) anthoxanthins, a group of colorless compounds further divided in several categories, including flavones, flavans, flavonols, flavanols, isoflavones, and their glycosides. Flavonols are mainly represented by myricetin, fisetin, quercetin and kaempferol.

2.3 Stilbenes

Stilbenes are structurally characterized by the presence of a 1,2-diphenylethylene nucleus with hydroxyls substituted on the aromatic rings, and exist in the form of monomers or oligomers. The best known compound is *trans*-resveratrol, possessing a trihydroxystilbene skeleton.

2.4 Tannins

Tannins are a group of water-soluble polyphenols having molecular weights from 500 to 3,000 which are subdivided into condensed and hydrolysable tannins, and commonly found complexed with alkaloids, polysaccharides and proteins, particularly the latter. On the basis of structural characteristics there are two groups, gallotannins and ellagitannins of hydrolysable tannins.

2.5 Diferuloylmethanes

Diferuloylmethanes are a small group of phenolic compounds with two aromatic rings substituted with hydroxyls, and linked by aliphatic chain containing carbonyl groups. There are also some other polyphenols such as hydroxytyrosol, a simple polyphenol presenting in olive fruits and olive oil [12,13].

Table 1. Classification and sources of dietary polyphenols

Class and subclass	Dietary polyphenol	Foods or beverages	Ref
Flavonoids			
Anthocyanidins	Cyanidin 3-galactoside	Fruits: blackberries, black currant, blueberries, black grape,	6
	Cyanidin 3-glucoside	elderberries, strawberries, cherries, plums, cranberry, pomegranate	14
	Cyanidin 3-arabinoside	juice, raspberry	15
	Cyanidin 3-xyloside	Others: red wine	16
	Malvidin		
	Delphinidin		
	Pelargonidin		
Anthoxanthins			
Flavonols	Myricetin	Vegetables: capers, celery, chives, onions, red onions, dock leaves,	7
	Fisetin	fennel, hot peppers, cherry tomatoes, spinach, sweet potato leaves,	17
	Quercetin	lettuce, celery, broccoli, Hartwort leaves, kale	14
	Kaempferol	Cereal: buckwheat, beans (green/yellow)	
	Isorhamnetin	Fruits: apples, apricots, grapes, plums, bilberries, blackberries, blueberries, cranberries, olive elderberries, currants, cherries, black currant juice, apple juice, ginkgo biloba	
		Spices and herbs: dill weed	
		Others: red wine, tea (green, black), tea (black beverage), cocoa powder, turnip (green), endive, leek	
Flavanones	Naringenin	Citrus fruits and juices: lemon, lemon juice, lime juice, orange,	18
	Eriodictyol	orange juice, grapefruit, tangerine juice	19
	Hesperetin	Spices and herbs: peppermint	20
Flavones	Apigenin	Fruits: celery, olives	14
	Luteolin	Vegetables: hot peppers, celery hearts, fresh parsley	21
		Spices and herbs: oregano, rosemary, dry parsley, thyme	22

Table 1. Cont.

Flavanols (Flavan-3-ols)	(+)-Catechin	Fruits: apples, apricots, grapes, peaches, nectarines, pears, plums, raisins, raspberries, cherries, blackberries, blueberries, cranberries Others: red wine, tea (green, black), chocolate (dark, milk), white wine, cocoa	23
	(-)-Epicatechin		24
	(-)-Epicatechin 3-gallate		
	Morin		
	(-)-Epigallocatechin		
	(-)-Epigallocatechin-3-gallate		
	(+)-Gallocatechin		
	Procyanidins		
	Prodelphinidins		
Isoflavones (Flavans)	Genistein	Fruits: grape seed/skin Others: soybean, soy nuts, soy flour/bread, tofu, miso, soy milk, tofu yogurt, soy cheese/sauce/hot dog	25
	Daidzein		
	Equol		
Flavonoid glycoside	Rutin	Fruits: lemon, orange, orange juice, grapefruit, tangerine juice	26
	Hesperidin		
	Naringin		
Phenolic acids Hydroxycinnamic acids	Caffeic acid	Fruits: blueberry, cranberry, pear, cherry(sweet), apple, orange, grapefruit, cherry juice, apple juice, lemon, peach, Vegetables: potato, lettuce, spinach Others: coffee beans, tea, coffee, cider	27
	Chlorogenic acid		
	Ferulic acid		
	Neochlorogenic acid		
	P-coumaric acid		
	Sinapic acid		
	Caftaric acids		
Hydroxybenzoic acids	Ellagic acid	Fruits: strawberry, raspberry grape juice(black/green), longan seed, pomegranate juice	28
	Gallic acid		29
	Corilagin		
Trihydroxy-stilbenes	Resveratrol	Fruits: grapes, peanuts, Others: red wine	30
	Trans-resveratrol		31
Tannins	Catechin polymers	Fruits: grape (dark/light) seed/skin, apple juice, strawberries, longan, raspberries, pomegranate, walnuts, muscadine grape, muscadine grape, peach, blackberry (juices/jams/jellies), olive, plum, Vegetables: chick pea, black-eyed peas, lentils, Cereal: haricot bean, Others: red wine, white wine, cocoa, chocolate, oak-aged red wine, tea, cider, tea, coffee, immature fruits	14
	Epicatechin polymers		29
	Ellagitannins		32
	Proanthocyanidins		
	Casuarictin		
	Sanguin H6		
Tannic acids			
Diferuloylmethane	Curcumin	herbal remedy, dietary spice turmeric	33


3. Bioactivities of dietary polyphenols

Oxidative stress is considered to play a pivotal role in the pathogenesis of aging and several degenerative diseases, such as atherosclerosis, cardiovascular disease, type II diabetes and cancer [1-3]. In order to cope with an excess of free radicals produced upon oxidative stress, humans have developed endogenous and exogenous mechanisms in order to maintain redox homeostasis. Among these, dietary polyphenols have been largely studied for their strong antioxidant capacities and other properties by which cell activities are regulated (Figures 2 and 3).

3.1 Antioxidant and free radical scavenging properties

In order to combat and neutralize the deleterious effects of ROS, various antioxidant strategies have evolved either by increasing the endogenous antioxidant enzyme defenses or by enhancing the non-enzymatic defenses through dietary or pharmacological means (Table 2). Dietary polyphenols have been reported to possess potent antioxidant activity by endogenous and exogenous mechanisms.

Figure 2. Bioactivities of dietary polyphenols.



Dihydrocaffeic acid was able to scavenge free radicals (superoxide anion, hydroxyl and peroxy radicals) in human EA.hy926 endothelial cells [42]. Curcumin and quercetin increased several antioxidant enzyme activities such as glutathione peroxidase (GPx), superoxide dismutase (SOD), catalase (CAT) or glutathione reductase (GR) *in vivo* and *in vitro* [8,9,44], and activated endogenous defense systems *in vitro* [40,45]. Hydroxytyrosol could increase CAT and SOD activities in rats fed a cholesterol-rich diet [35].

The transcription factor Nrf2 regulates the basal and inducible expression of numerous detoxifying and antioxidant genes. The Nrf2–Kelch-like ECH-associated protein 1 (Keap1)-ARE system is now recognized as one of the major cellular defence mechanisms against oxidative and xenobiotic stresses [46]. (-)-Epigallocatechin gallate (EGCG) and (-)-epicatechin gallate (ECG) induced ARE-mediated gene expression through the activation of MAPK proteins (ERK, JNK and p38) in HepG2-ARE-C8 cell [10]. Tanigawa *et al.* reported that quercetin-induced ARE activity involves upregulation of Nrf2 through the regulation of both transcription and posttranscription sites and repression of Keap1 by affecting the posttranscription site in HepG2 cells [48]. Curcumin could increase the expression of glutathione S-transferase P1 (GSTP1) by activating ARE and Nrf2 in HepG2 cells [40].

Figure 3. Mechanisms of the biological effects of dietary polyphenols.




Table 2. Antioxidant and free radical scavenging properties of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Epigallocatechin, EGCG, ECG	Inhibiting lipoxygenase and cyclooxygenase	In human colon mucosa and colon tumor tissues	<i>In vitro</i>	34
EGCG ECG	Inducing ARE-mediated gene expression through the activation of MAPK proteins (ERK, JNK and p38)	In HepG2-ARE-C8 cell	<i>In vitro</i>	10
Hydroxytyrosol	Increasing CAT and SOD activities	In rats fed a cholesterol-rich diet	<i>In vivo</i>	35
	Inhibiting the activities of 12-lipoxygenase and 5-lipoxygenase	In rat platelets and rat polymorphonuclear leukocytes (PMNL)	<i>In vitro</i>	36
	Reducing leukotriene B4 production			

Table 2. Cont.

Catechin Proanthocyanidin B4	Increasing CAT, glutathione S-transferase (GST) and SOD activities Elevating cellular GSH content	In cardiac H9C2 cells	<i>In vitro</i>	37
Curcumin	Inhibiting CYP1A2, CYP3A4, CYP2B6, CYP2D6, and CYP2C9	The plasmids with human cytochrome P450 NADPH reductase	<i>In vitro</i>	38
	Inhibiting mitochondrial proton F0F1-ATPase/ATP synthase	Rat brain F0F1-ATPase	<i>In vitro</i>	39
	Increasing the expression of GSTP1 by activating ARE and Nrf2	In HepG2 cells	<i>In vivo</i>	40
	Increasing CAT, SOD activity and heat shock proteins 70 expression Decreasing the activity of iNOS Decreasing malondialdehyde (MDA), NO(2)(-) + NO(3)(-) and myeloperoxidase (MPO) level and serum transaminase concentration	In rat model	<i>In vivo</i>	8
Kaempferol-3-O-galactoside	Inhibiting human recombinant synovial phospholipase A2 (PLA2)	In mice	<i>In vivo</i>	41
EGCG, Quercetin, Kaempferol Morin, Apigenin, Daidzein, ECG	Inhibiting mitochondrial proton F0F1-ATPase/ATP synthase	Rat brain F0F1-ATPase	<i>In vitro</i>	39
Ellagic acid Gallic acid Corilagin	Inhibiting tyrosinase, xanthine oxidase, and the formation of superoxide radical	In substrate of L-tyrosine	<i>In vitro</i>	29
Dihydrocaffeic acid	Enhancing eNOS activity and protein expression Scavenging intracellular ROS	In human EA.hy926 endothelial cells	<i>In vitro</i>	42
Caffeic acid (+)-catechin	Inhibiting peroxynitrite-mediated oxidation of dopamine	In dopamine	<i>In vitro</i>	43
Quercetin	Preventing lactate dehydrogenase (LDH) leakage Increasing SOD, CAT, GSH, GPx, and GR activity	In mouse liver	<i>In vivo</i>	9
	Decreasing MDA and lipoperoxidation Increasing Cu/Zn SOD and GPx mRNA	In HepG2 cells	<i>In vitro</i>	44
	Increasing the expression and activity of NADPH:quinone oxidoreductase-1 (NQO1)	In the MCF-7 human breast carcinoma cellse	<i>In vitro</i>	45
	Enhancing γ -glutamylcysteine synthetase (γ -GCS)	In HepG2 cells	<i>In vitro</i>	47
	Enhancing the ARE binding activity and Nrf2-mediated transcription activity Upregulating and stabilizing Nrf2 Reducing the level of Keap1 protein	In HepG2 cells	<i>In vitro</i>	48

Table 2. Cont.

Resveratrol	Inhibiting O-acetyltransferase and sulfotransferase activities Preventing the oxidative DNA damage	In male Wistar rats treated with potassium bromate	<i>In vivo</i>	49
	Inhibiting the production of H ₂ O ₂ and MPO activity Increasing GSH levels and SOD activities Decreasing the levels of MPO and oxidized GR	In mouse skin	<i>Ex vivo</i>	50
	Reducing PhIP-DNA-adduct formation by O-acetyltransferase and sulfotransferase catalysis	In primary cultures of human mammary epithelial cells	<i>In vitro</i>	51
	Inhibiting the expression and activity of CYP 1A1/1A2	In microsomes and intact HepG2 cells	<i>In vitro</i>	52
	Inhibiting mitochondrial proton F0F1-ATPase/ATP synthase	Rat brain and liver F0F1-ATPase	<i>In vitro</i>	39
	Suppressing CYP1A1 and IL-1 β transcription by blocking aryl hydrocarbon receptor		<i>Ex vivo</i> <i>In vivo</i>	53
	(-)-Epicatechin Procyanidin EGCG, ECG	Inhibiting recombinant human platelet 12-lipoxygenase and 15-lipoxygenase	In rabbit smooth muscle cells and in J774A.1 cells	<i>In vitro</i>

3.2. Anti-atherosclerosis and cardioprotection

Studies have shown that some of dietary polyphenols exerted anti-atherosclerosis and cardioprotection (Table 3). Oleuropein inhibited the oxidation of low density lipoprotein (LDL) *in vitro* [61]. Quercetin decreased lipid peroxidation, upregulated the expression of serum high density lipoprotein (HDL)-associated paraoxonase 1(PON-1) in the HuH7 human hepatoma cell line [66], inhibited oxidized LDL (oxLDL)-triggered apoptosis, and increased intracellular glutathione (GSH) downregulation in COS-1 cells [68].

Proanthocyanidin could significantly reduce cardiomyocyte apoptosis by inhibiting ischemia/reperfusion-induced activation of JNK-1 and c-Jun in Male Sprague Dawley rats [74]. Furthermore, proanthocyanidin could regulate the levels of CD36 mRNA and protein in oxLDL treated peripheral blood mononuclear cells [73]. Resveratrol showed that *in vitro* it could decrease the expression of vascular cell adhesion molecule-1 (VCAM-1) [64], cyclooxygenase-2 (COX-2) [55], and matrix metalloproteinase-9 (MMP-9) mRNA [56] through suppression of activation of nuclear factor AP-1 [55]. Hydroxytyrosol could not only lower serum total cholesterol (TC) and low density lipoprotein cholesterol (LDL-C), but also slow the lipid peroxidation process in rats fed a cholesterol-rich diet [35].

Table 3. Anti-atherosclerosis and cardioprotection of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Resveratrol	Suppressing the expression and activity of COX-2 Suppressing activation of AP-1	In human mammary epithelial cells	<i>In vitro</i>	55
	Inhibiting the activity and expression of MMP-9	In U937 cells	<i>In vitro</i>	56
	Enhancing myocardial angiogenesis by induction of VEGF, thioredoxin-1 (Trx-1), and HO-1	In male Sprague Dawley rats	<i>In vivo</i>	57
	Inhibiting the expression and binding activity of the monocyte chemotactic protein-1 (MCP-1) receptor, CC-chemokine receptor-2 (CCR2)	on THP-1 monocytes	<i>In vitro</i>	58
	Increasing NO and NOS levels Increasing intracellular cyclic GMP (cGMP) level and decreasing atrial natriuretic peptide (ANP) and brain natriuretic peptide (BNP) levels	In cultured rat cardiac fibroblasts	<i>In vitro</i>	59
(-)-Epicatechin	Inhibiting 7 β -OH-cholesterol formation	In endothelial cells	<i>In vitro</i>	60
Hydroxytyrosol	Preventing platelet aggregation and eicosanoid formation Inhibiting thromboxane B2 production	In platelet rich plasma	<i>In vitro</i>	61
	Inhibit thromboxane B2 production	In patients with uncomplicated type I diabetes	<i>In vivo</i>	62
Hydroxytyrosol Oleuropein Caffeic acid	Inhibiting leukotriene B4 generation Inhibiting 5-lipoxygenase	In rat peritoneal leukocytes	<i>In vitro</i>	63
Oleuropein Hydroxytyrosol Resveratrol	Reducing monocytoïd cell adhesion to stimulated endothelium Decreasing VCAM-1 mRNA and protein	In human umbilical vein endothelial cells (HUVECs)	<i>In vitro</i>	64
Oleuropein	Decreasing creatine kinase and GSH release	In the isolated rat heart	<i>Ex vivo</i>	65
Quercetin	Upregulating the expression of serum HDL-associated PON-1	In the HuH7 human hepatoma cell line	<i>In vitro</i>	66
Kaempferol Apigenin	Inducing interferon-gamma (IFN- γ) gene expression Downregulating IL-4 gene expression	In peripheral blood mononuclear cells	<i>In vitro</i>	67
	Increasing the intracellular GSH and activating γ -GCS heavy subunit (GCS(h)) promoter	In COS-1 cells	<i>In vitro</i>	68
EGCG and ECG	Inhibiting rat VSMCs adhesion on collagen and laminin Interference with VSMC's integrin β 1 receptor and binding to extracellular matrix (ECM) proteins	In rat VSMCs	<i>In vitro</i>	69
Genistein	Decreasing hydroxyproline concentrations Suppressing the progression of myocardial fibrosis	In Long-Evans Tokushima Otsuka non-diabetic rats	<i>In vivo</i>	70
Genistein Daidzein	Incorporating into LDLs, increasing their oxidation resistance and antiproliferative efficacy	In cultured U937 cells	<i>Ex vivo</i>	71

Table 3. Cont.

Procyanidins	Decreasing leukotriene-prostacyclin ratio in plasma	In humans and human aortic endothelial cells	<i>In vivo</i> <i>In vitro</i>	72
Proanthocyanidin	Inhibiting CD36 mRNA expression	In peripheral blood mononuclear cell	<i>In vitro</i>	73
Proanthocyanidin	Reducing cardiomyocyte apoptosis by inhibiting ischemia–reperfusion-induced activation of JNK-1 and c-Jun	In Male Sprague Dawley rats	<i>In vivo</i>	74
Hydroxytyrosol	Lowering serum TC and LDL-C Slowing the lipid peroxidation process	In rats fed a cholesterol-rich diet	<i>In vivo</i>	35

3.3 Neuroprotective effects on anti-aging and neurodegenerative diseases

Recently, there has been considerable interest in the neuroprotective effects of dietary polyphenols (Table 4), especially in the context of their modes of action as antioxidants [6]. Resveratrol had an impact on cognitive deficits by activating the phosphorylation of protein kinase C (PKC), secreting transthyretin to prevent A β aggregation in cultured rat hippocampal cells [77], and stimulating AMP kinase activity in Neuro2a cells and primary neurons [75]. EGCG stimulated the deacetylase activity of recombinant silent information regulator two ortholog 1 (SIRT1) protein in human HT29 cells [80]. Curcumin could disrupt existing plaques and restore distorted neurites in an Alzheimer mouse model [84]. They had been considered as therapeutic agents for altering brain aging processes, and as possible neuroprotective agents in progressive neurodegenerative disorders such as Parkinson's and Alzheimer's diseases.

Table 4. Neuroprotective effects of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Hydroxytyrosol	Attenuating Fe ²⁺ - and NO-induced cytotoxicity Increasing cellular ATP Reducing lipid peroxidation Hyperpolarizing basal mitochondrial membrane potential	In murine-dissociated brain cells and mice	<i>In vitro</i> <i>Ex vivo</i>	12
Resveratrol	Stimulating AMP kinase activity	In Neuro2a cells and primary neurons	<i>In vitro</i>	75
	Preventing fibrosis, NF- κ B activation and TGF- β increases induced by chronic CCl(4) treatment	In rats	<i>In vivo</i>	76
	Activating the phosphorylation of PKC Secreting transthyretin to prevent A β aggregation	In cultured rat hippocampal cells	<i>In vitro</i>	77
	Protecting dopaminergic neurons Activating sirtuin family of NAD-dependent histone deacetylases	In organotypic midbrain slice culture	<i>In vitro</i>	78

Table 4. Cont.

EGCG ECG Myricetin	Inhibiting IL-6, IL-8, VEGF and prostaglandin E2 (PGE2) production Attenuating the expression of COX-2 and activation of NF- κ B Inducing the expression of MAPK phosphatase-1 Suppressing the phosphorylation of MAPK (p38 and JNK)	In human astrocytoma U373MG cells	<i>In vitro</i>	79
	Attenuating disruption of mitochondrial membrane potential and release of cytochrome c Decreasing the activities of caspase-9 and caspase-3 and increase in the Bax to Bcl-2 ratio	In rat PC12 cells	<i>In vitro</i>	115
	Stimulating the deacetylase activity of recombinant SIRT1 protein	In human HT29 cells	<i>In vitro</i>	80
	Increasing the activities of PKC and ERK1/2 Decreasing the expression of Bax, Bad, and Mdm2 Increasing the expression of Bcl-2, Bcl-w, and Bcl-xL	In human neuroblastoma SH-SY5Y cell	<i>In vitro</i>	81
Catechin Quercetin Genestein Naringenin	Attenuating the apoptotic injury induced <i>N</i> -methyl-4-phenyl-1,2,3,6-tetrahydropyridinium hydrochloride (MPP+)	In mesencephalic dopamine neurones	<i>In vivo</i>	82
Epicatechin Kaempferol	Protecting neurons from oxLDL-induced apoptosis by inhibiting the activation of JNK, c-Jun and caspase-3	In cultured primary neurons	<i>In vitro</i>	83
Curcumin	Disrupting existing plaques and restoring distorted neurites Crossing the blood-brain barrier and labels senile plaques and cerebrovascular amyloid angiopathy	In an Alzheimer mouse model In APP ^{swe} /PS1 ^{dE9} mice	<i>In vivo</i>	84

3.4 Anti-inflammatory properties

Oxidative stress induced inflammation is mediated by the activation of NF- κ B and AP-1. It affects a wide variety of cellular signaling processes leading to generation of inflammatory mediators and chromatin remodeling [95,96]. The latter allows expression of pro-inflammatory genes such as interleukin-1 β (IL-1 β), IL-8, tumor necrotic factor alpha (TNF- α), and inducible nitric oxide synthase (iNOS). The undesired effects of oxidative stress have been found to be controlled by the antioxidant and/or anti-inflammatory effects of dietary polyphenols such as curcumin and resveratrol *in vivo* and *in vitro* [88-90,95,97] (Table 5). Resveratrol inhibited pro-inflammatory gene expression via inhibition of inhibitory κ B (I κ B), thus inhibiting NF- κ B transactivation, as well as restoring transrepressive pathways through the activation of histone deacetylases in RAW 264.7 cells [89].

Table 5. Anti-inflammatory effects of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Procyanidins	Inhibiting transcription and secretion of IL-1 β	In peripheral blood mononuclear cells	<i>In vitro</i>	85
EGCG ECG	Inducing apoptosis by activating caspases 3, 8, and 9	In Isolated peripheral blood monocytes	<i>In vitro</i>	86
	Downregulating CD11b expression Attenuating adhesion and migration of peripheral blood CD8+T cells	In peripheral blood CD8+ T cells	<i>In vitro</i>	87
Resveratrol	Inhibiting stimulation of caspase-3 and cleavage of PARP induced by IL-1 β	In human articular chondrocytes	<i>In vitro</i>	88
	Suppressing the expression of iNOS mRNA and protein by inhibiting the activation of NF- κ B Inhibiting NO generation	In RAW 264.7 cells	<i>In vitro</i>	89
	Upregulating MAP kinase phosphatase-5	In prostate cells	<i>In vitro</i>	90
Apigenin	Blocking the expression of intercellular adhesion molecule-1 (ICAM-1), VCAM-1, and E-selectin Inhibiting prostaglandin synthesis and IL-6, 8 production	In human endothelial cells	<i>In vitro</i>	91
Luteolin	Inhibiting the upregulation of THP-1 adhesion and VCAM-1 expression	In HUVECs	<i>In vitro</i>	92
Quercetin	Inhibiting the activity of the NF- κ B			
	Inhibiting NO production and iNOS protein expression	In NR8383 macrophages	<i>In vitro</i>	93
Anthocyanins	Localizing into endothelial cells	In human microvascular endothelial cells	<i>In vitro</i>	94
Hydroxy-cinnamic acids	Reducing the upregulation of IL-8, MCP-1, and ICAM-1			
Curcumin	Decreasing MPO activity and TNF- α on chronic colitis Reducing nitrites levels and the activation of p38 MAPK Downregulating COX-2 and iNOS expression	In rats	<i>In vivo</i>	95
	Upregulating MAP kinase phosphatase-5	In prostate cells	<i>In vitro</i>	90
	Suppressing the induction of COX-2 and iNOS Inhibiting the expression of ICAM-1 and MCP-1 Suppressing the Janus kinase (JAK)-STAT via activation of Src homology 2 domain-containing protein tyrosine phosphatases (SHP-2)	In both rat primary microglia and murine BV2 microglial cells	<i>In vitro</i>	97

On the other hand, to counter the effects of oxidative stress, the cells also concomitantly express protective antioxidants such as glutamate cysteine ligase (GCL), manganese superoxide dismutase

(MnSOD), and heme oxygenase-1(HO-1). In addition, expression of these antioxidant genes via modulation of MAPK-ARE-Nrf2 pathway is upregulated by EGCG and ECG in HepG2-ARE-C8 cell [10]. Apigenin, luteolin and quercetin had also been reported to inhibit inflammatory responses by downregulating the expression of iNOS and adhesion molecules in NR8383 macrophages and human endothelial cells [91-93].

3.5 Antimutagenic/anticarcinogenic properties

Dietary polyphenols could modulate diverse biochemical processes involved in carcinogenesis (Table 6). Curcumin exerted antitumor activities by inhibition of cellular proliferation and angiogenesis, blockade of tumor cell cycle progression, and induction of programmed cell death in vivo and in vitro [109,110]. Cellular signaling cascades mediated by NF- κ B or AP-1 acted as a centerplay in regulating many of aforementioned biochemical processes [102,110].

Table 6. Antimutagenic/anticarcinogenic properties of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Hydroxytyrosol	Inhibiting cell proliferation Inducing apoptosis by arresting the cells in the G0/G1 phase with a concomitant decrease in the cell percentage in the S and G2/M phases	In human promyelocytic leukaemia cells HL60	<i>In vitro</i>	98
Resveratrol	Inhibiting cell proliferation and downregulating telomerase activity	In human colon tumor cells	<i>In vitro</i>	99
	Inducing apoptosis mediated by p53-dependent pathway	In HepG2 cells	<i>In vitro</i>	100
	Inhibiting cell proliferation by interfering with an estrogen receptor- α (ER α)-associated PI3K pathway	In estrogen-responsive MCF-7 human breast cancer cells	<i>In vitro</i>	101
	Suppressing COX-2 expression by blocking the activation of MAPKs and AP-1	In dorsal skin of female ICR mice	<i>In vitro</i>	102
	Decreasing the expression of COX-1, COX-2, c-myc, c-fos, c-jun, transforming growth factor-beta1 (TGF- β 1) and TNF- α	In mouse skin	<i>Ex vivo</i>	50
	Inhibiting oncogenic disease through the inhibition of protein kinase CKII activity	In HeLa cell lysates	<i>In vitro</i>	103
	Inhibiting the Ca(2+)-dependent activities of PKC α and PKC β I	On the activities of PKC isozymes	<i>In vitro</i>	104
	Inhibiting nitrobenzene(NB)-DNA adducts and NB-Hb adducts	In male Kunming mice	<i>In vivo</i>	105
Chlorogenic acid	Inhibiting the formation of DNA single strand breaks	In supercoiled pBR322 DNA	<i>In vitro</i>	106
Quercetin Luteolin	Blocking EGFR tyrosine kinase activity	In MiaPaCa-2 cancer cells	<i>In vitro</i>	107

Table 6. Cont.

Myricetin Apigenin Quercetin Kaempferol	Inhibiting human CYP1A1 activities Inhibiting the formation of diolepoxide 2(DE2) and B[a]P activation	On 7-ethoxyresorufin O-deethylation	<i>In vitro</i>	26
Silymarin Hesperetin Quercetin Daidzein	Interacting with P-glycoprotein and modulating the activity of ATP-binding cassette transporter, breast cancer resistance protein (BCRP/ABCG2)	In two separate BCRP-overexpressing cell lines	<i>In vitro</i>	108
EGCG	Inhibiting telomerase	In human cancer cells In nude mice models	<i>In vitro</i> <i>In vivo</i>	114
Curcumin	Suppressing proliferation and angiogenesis Inhibiting NF- κ B-regulated gene products (cyclin D1, c-myc, Bcl-2, Bcl-xL, cellular inhibitor of apoptosis protein-1, COX-2, MMP, and VEGF)	In various pancreatic cancer cell lines and nude mice	<i>In vitro</i> <i>In vivo</i>	109
	Inducing apoptosis by sustained phosphorylation of JNK and p38 MAPK Inhibiting NF- κ B transcriptional activity Inducing phosphorylation of c-jun and stimulation of AP-1 transcriptional activity	In HCT116 cells	<i>In vitro</i>	110
	Inducing apoptosis through activation of caspase-8, BID cleavage and cytochrome c release Suppressing ectopic expression of Bcl-2 and Bcl-xl	In human acute myelogenous leukemia HL-60 cells	<i>In vitro</i>	111
	Inhibiting the Akt/mTOR/p70S6K pathway and activating the ERK1/2 pathway Inhibiting tumor growth and inducing autophagy	In U87-MG and U373-MG malignant glioma cells In the subcutaneous xenograft model of U87-MG cells	<i>In vitro</i> <i>In vivo</i>	112

Resveratrol could block the activation of MAPKs and AP-1 in the skin of mice [102]. Consumption of berries and red fruits rich in polyphenols contributed to the reduction of cancer through many mechanisms such as *in vitro* inhibiting human cytochrome P450-dependent monooxygenases 1A1 (CYP1A1) activities [26], blocking the epidermal growth factor receptor (EGFR) tyrosine kinase activity [107], and decreasing protein kinase CKII activity [103].

3.6 Maintenance of gastrointestinal health and effects on digestive enzymes

It had been reported that digestive enzymes such as lipase, α -amylase, and α -glucosidase, were inhibited by proanthocyanidins and tannins in young chicks, which decreased the digestibility of protein, starch and lipid [119, 120]. Resveratrol could inhibit pancreatic bile salt-dependent lipase (BSDL) activity, expression and secretion in the rat pancreatic AR4-2J cells [121]. Cyanidin-3 α -O-rhamnoside and quercetin-3 α -O-rhamnoside could inhibit α -glucosidase and advanced glycation end product (AGE) formation *in vitro* [123]. The inhibition of digestive enzymes by dietary polyphenols

may represent an under-reported mechanism for delivering some of the health benefits attributed to a diet rich in fruit and vegetables.

3.7 Modulation of signal transduction pathways

Table 7. Effects of dietary polyphenols on signal transduction pathways.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Curcumin	Inhibiting both myeloid differential factor 88 (MyD88)-and TIR domain-containing adapter inducing IFN- β (TRIF)-dependent pathways Inhibiting homodimerization of Toll-like receptor 4(TLR4) Suppressing the activation of NF- κ B by inhibiting I κ B kinase β activity in MyD88-dependent pathway Inhibiting IFN-regulatory factor 3 (IRF3) activation	In 293T cells	<i>In vitro</i>	124
	Inhibiting the level of NOS mRNA and protein Suppressing NF- κ B activation through inhibitory of I κ B kinase activity	In macrophages	<i>In vitro</i>	125
	Suppressing COX-2 expression by inhibiting AP-1 and NF- κ B	In BV2 microglial cells	<i>In vitro</i>	126
	Inhibiting IL-6-inducible STAT3 phosphorylation and nuclear translocation	In human multiple myeloma cells	<i>In vitro</i>	118
	Upregulating CYP3A4 via pregnane X receptor (PXR) activation Activating the electrophile responsive element (EpRE) of HO-1 and enhancing the gastrointestinal (GI)-GPx activity	In HepG2 cells	<i>In vitro</i>	127
	Suppressing JAK-STAT inflammatory signaling through activation of SHP-2	In both rat primary microglia and murine BV2 microglial cells	<i>In vitro</i>	97
	Proanthocyanidins	Promoting apoptosis through alterations in Cdk1-Cdk-cyclin cascade, and caspase-3 activation via loss of mitochondrial membrane potential	In human epidermoid carcinoma A431 cells	<i>In vitro</i>
Proanthocyanidins	Inhibiting the phosphorylation of ERK1/2, JNK and p38 Inhibiting the activation of NF- κ B/p65 through inhibition of degradation of I κ B α and activation of I κ B kinase α	In SKH-1 hairless mice	<i>In vivo</i>	129

Table 7. Effects

Caffeic acid	Modulating ceramide-induced signal transduction pathway and NF- κ B activation Inhibiting protein tyrosine kinase activity	In U937 cells	In vitro	113
Quercetin	Inhibiting phosphorylation of JNK and p38 MAPK on ROS-mediated signaling	In HUVECs	In vitro	117
	Modulating Akt/PKB and ERK1/2 signalling cascades on neuronal viability	In primary cortical neurons	In vitro In vivo	130
Equol	Mediating rapid vascular relaxation by Ca ²⁺ -independent activation of eNOS/Hsp90 involving ERK1/2 and Akt phosphorylation	In human endothelial cells	In vitro	131
Resveratrol	Inhibiting monocyte CCR2 binding activity in an NO-, MAPK- and PI3K-dependent manner Inhibiting CCR2 mRNA in an NO- and MAPK-independent, PI3K-dependent manner	on THP-1 monocytes	In vitro	58
	Inhibiting proliferation of cardiac fibroblasts by NO-cGMP signaling pathway	In cultured rat cardiac fibroblasts	In vitro	59
	Inducing phase II genes by regulating ARE/EpRE activation Modifying the capability of Keap1 in sequestering Nrf2	In PC12 cells	In vitro	132

Dietary polyphenols may not merely exert their diverse biological effects as free radical scavengers, but may also modulate cellular signaling processes by affecting signal transduction pathways [122] (Table 7). Studies have been reported that curcumin could in vitro modulate NF- κ B activation [124], AP-1 DNA binding [126], signal transducer and activator of transcription-3 (STAT3) phosphorylation [118]. Resveratrol exerted protection in vitro through PI3-kinase/Akt pathway, MAPK proteins (ERK, JNK and P38) activation [58], and the translocation into the nucleus of Nrf2 [132]. Resveratrol could also upregulate the expressions of GCL, MnSOD, and HO-1 against oxidative stress via MAPK-ARE-Nrf2 pathway in PC12 cells [132].

3.8 Improvement of endothelium functions

Several studies have indicated that red wine polyphenolic compounds (RWPCs) were able to inhibit proliferation and migration of vascular cells (Table 8). RWPCs induced nitric oxide (NO)-mediated endothelium-dependent relaxations in isolated arteries. The activation of endothelial NO synthase (eNOS) was due to two distinct mechanisms: (a) an increase in [Ca²⁺]_i and (b) a phosphorylation of eNOS by the PI3-kinase/Akt pathway [137]. In addition, RWPCs caused endothelium-derived hyperpolarizing factor (EDHF)-mediated relaxations of isolated arteries consecutively to a localized and controlled formation of superoxide anions leading to the activation of the PI3-kinase/Akt pathway [136]. RWPCs also increased endothelial prostacyclin release and inhibited the synthesis and the effects of endothelin-1 in endothelial cell [139,141].

Table 8. Protective effects of dietary polyphenols on endothelial cells and blood vessels

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
EGCG Quercetin	Inhibiting apoptosis through modulation of Bcl-2 and Bax Inhibiting nuclear transactivation of p53 Decreasing the activity of caspase-3 Blocking JNK- and p38 MAPK-related signaling	In HUVECs	<i>In vitro</i>	117
RWPCs	Inhibiting the expression of VEGF mRNA and protein Preventing the activation of the p38 MAPK pathway	In VSMCs	<i>In vitro</i>	133
	Inhibiting the invasion and migration of VSMCs Inhibiting pro-MMP-2 expression and its activation via inhibition of membrane type 1-MMP (MT1-MMP) activity	In VSMCs	<i>In vitro</i>	134
	Inhibiting VSMCs migration through inhibiting the PI3K activity and p38 MAPK phosphorylation Inhibiting the phosphorylation of MKK3/6	In cultured VSMCs	<i>In vitro</i>	135
	Inducing EDHF-mediated relaxations through activation of the PI3-kinase/Akt pathway	In porcine coronary arteries	<i>In vivo</i>	136
	Increasing intracellular Ca ²⁺ and activate tyrosine kinases Increasing NO production	In bovine aortic endothelial cells	<i>In vitro</i>	137
	Inhibiting NADPH oxidase activity and/or reducing endothelin-1(ET-1) release	In Twelve-week-old male Wistar rats	<i>In vivo</i>	138
	Inhibiting the synthesis of ET-1	In cultured bovine aortic endothelial cells	<i>In vitro</i>	139
	Elevating NO and prostacyclin (PGI ₂)	In rats	<i>In vivo</i>	140
	Enhancing PGI ₂ release	In endothelial cell	<i>In vitro</i>	141
Cy3G	Enhancing eNOS activity and expression Inducing NO production Regulating phosphorylation of eNOS and Akt Increasing cGMP production	In bovine vascular endothelial cells	<i>In vitro</i>	142
EGCG	Having endothelial-dependent vasodilator actions Activating phosphatidylinositol 3-kinase, Akt, and eNOS	In bovine aortic endothelial cells	<i>In vitro</i>	143
	Increasing eNOS activity Inducing a sustained activation of Akt, ERK1/2, and eNOS Ser1179 phosphorylation	In bovine aortic endothelial cells	<i>In vitro</i>	144
Catechins	Reducing the vascularization induced by the angiogenin-like protein on chicken CAM	In chichen	<i>In vitro</i>	145
Activin	Reducing ICAM-1, VCAM-1 and E-selectin	In systemic sclerosis	<i>In vivo</i>	146
Proanthocyanidin	Downregulating VCAM-1 expression; Decreasing TNF α -induced adherence of T-cells to HUVECs	In primary HUVECs	<i>In vitro</i>	147
Procyanidins Flavan-3-ols	Inhibiting angiotensin I converting enzyme (ACE) activity	In two substrates	<i>In vitro</i>	148

RWPCs could prevent matrix metalloproteinases-2 (MMP-2) activation and vascular endothelial growth factor (VEGF) expression in vascular smooth muscle cells (VSMCs) [133,134]. All these mechanisms might contribute to explain the vasodilatory, vasoprotective and anti-hypertensive effects of polyphenols *in vivo*.

Cyanidin-3-glucoside (Cy3G) and EGCG could enhance vascular eNOS activity and improve vascular endothelial function in bovine vascular endothelial cells [142]. Catechins had anti-angiogenic effects by reducing the vascularization on the chicken chorioallantoic membrane (CAM) [145].

3.9 Protective effect on immune cell functions

Dietary polyphenols appear to have a protective effect on immune cell functions. Alvarez *et al.* showed that leukocyte functions were improved in prematurely aging mice after five weeks of diet supplementation with polyphenol-rich cereals [149]. They could increase macrophage chemotaxis, phagocytosis, microbicidal activity, and natural killer function, and increase lymphoproliferation and IL-2 release in response to concanavalin A and lipopolysaccharide.

Curcumin could prevent tumor-induced T cell apoptosis by downregulating Bax level and augmenting Bcl-2 expression and restore cytokine-dependent Jak-3/Stat-5a signaling pathway in T cells of tumor bearer [150]. Caffeic acid, ellagic acid, and ferulic acid could inhibit apoptosis through the Bcl-2 independent mechanism in normal human peripheral blood mononuclear cells [116]. Thus, regular intake of these compounds will protect and improve quality of life.

3.10 Antiallergic activity

The incidence of type I allergic disorders have been increasing worldwide, particularly, the hypersensitivity to food. Akiyama and his coworkers reported that the apple condensed tannins intake would inhibit the development of the oral sensitization, and the inhibition could correlate with the rise in the population of TCR $\gamma\delta$ -T cells in the intestinal intraepithelial lymphocytes [151]. Moreover, the apple condensed tannins could inhibit the release of histamine from rat basophilic leukemia (RBL-2H3) cells stimulated by the antigen-stimulation and from rat peritoneal mast cells stimulated by compound 48/80. They also inhibited hyaluronidase activity and increase in intracellular free calcium concentration in RBL-2H3 cells stimulated with the antigen [152].

3.11 Antidiabetic effects

Johnston and coworkers demonstrated that glucose uptake into cells under sodium-dependent conditions was inhibited by flavonoid glycosides and non-glycosylated polyphenols in polarised Caco-2 intestinal cells [154]. Under sodium-free conditions, aglycones and non-glycosylated polyphenols inhibited glucose uptake whereas glycosides and phenolic acids were ineffective. These data suggest that aglycones inhibit facilitated glucose uptake whereas glycosides inhibit the active transport of glucose. The non-glycosylated dietary polyphenols appeared to exert their effects via steric hindrance, while EGCG, ECG and (-)-epigallocatechin were effective against both transporters.

More recently, Koboyashi *et al.* have shown that the green tea polyphenols EGCG and ECG also inhibited glucose transport, possibly by sodium-dependent glucose transporter 1 (SGLT1) inhibition in

the rabbit small intestine [155]. Song et al have presented evidence for quercetin-mediated inhibition of the facilitated diffusion glucose transporter 2 (GLUT2) in Chinese hamster ovary cells [156].

Anthocyanins inhibited α -glucosidase activity and reduced blood glucose levels after starch-rich meals. This is a proven clinical therapy for controlling type II diabetes [158] (Table 9).

Table 9. Antidiabetic activity of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Curcumin	Inhibiting diabetes-induced elevation in the levels of IL-1 β , VEGF, and NF- κ B Decreasing oxidatively modified DNA and nitrotyrosine	In streptozotocin-induced diabetic rats	<i>In vivo</i>	153
EGCG, ECG, (-)-epigallocatechin	Inhibiting SGLT1 and sodium-free GLUT	In polarised Caco-2 intestinal cells	<i>In vitro</i>	154
	Inhibiting SGLT1 and glucose uptake	In the rabbit small intestine	<i>In vivo</i>	155
Quercetin	Reducing blood glucose levels Inhibiting sodium-dependent vitamin C transporter 1 (SVCT1) and GLUT2	In Chinese hamster ovary cells	<i>In vitro</i>	156
Mangiferin	Inhibiting sucrase, isomaltase, and aldose reductase	In rats	<i>In vivo</i>	157
Tannins Anthocyanin	Inhibiting α -amylase and α -glucosidase	In the substrate of 2-chloro-4-nitro-phenyl-4-O-b-D-galactopyranosyl-maltoside	<i>In vitro</i>	158

3.12 Regulation of cell cycle progression

It was demonstrated that resveratrol and proanthocyanidins could regulate cell cycle progression by upregulating p21 expression, G1 phase arrest and downregulating cyclin D1/D2–Cdk6 *in vitro* [163–165, 170] (Table 10).

3.13 Modulation of hormonal effects and contraceptive activity

Some studies showed that dietary polyphenols could modulate the level of hormone. Resveratrol could exert mixed estrogen agonist/antagonist activities in mammary tumor models. It could affect the expression of 17 β -estradiol-responsive progesterone receptor (PR) and presnelin 2 proteins *in vitro* and *in vivo* [159]. Bhat *et al.* showed that resveratrol exhibited antiestrogenic properties and inhibited the levels and activity of PR by downregulating α (1)-integrin expression in human endometrial adenocarcinoma cells [160].

Otake and his coworkers demonstrated that quercetin and resveratrol potently reduced estrogen sulfotransferase (EST) activity and inhibited sulfation of 17 β -estradiol in normal human mammary epithelial cells [161]. Both of the compounds potently inhibited recombinant human EST. In fact, they could serve as substrates for EST. Gossypol, a polyphenolic compound from cotton seed, had contraceptive activity and could inhibit 11 β -hydroxysteroid dehydrogenase and cause hypokalemia in some men [162].

Table 10. Regulate cell cycle progression of dietary polyphenols.

Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Resveratrol	Upregulating p21 expression and cause G1 phase arrest	In HepG2 cells	<i>In vitro</i>	163
	Inhibiting cyclin D1/D2-cdk6, cyclin D1/D2-cdk4, and cyclin E-cdk2 complexes	In human epidermoid carcinoma A431 cells	<i>In vitro</i>	164
	Downregulating cyclin D1/Cdk4 complex and Upregulating cyclin E and A expression	In the human colonic adenocarcinoma cell line Caco-2	<i>In vitro</i>	165
	Decreasing in the hyperphosphorylated form of pRb and increasing in hypophosphorylated pRb Downregulating the protein expression of E2F (1-5) family members of transcription factors and their heterodimeric partners DP1 and DP2 Leading to a G0/G1 arrest	In human epidermoid carcinoma A431 cells	<i>In vitro</i>	166
	Inhibiting the expression of cyclin B1, D1, A1 and β -catenin	In six human cancer cell lines (MCF7, SW480, HCE7, Seg-1, Bic-1, and HL60)	<i>In vitro</i>	167
	Arresting cell cycle in the G1-S phase	In VSMCs	<i>In vitro</i>	168
	Upregulating the expression of cyclins A, E, and B1	In human SK-Mel-28 melanoma cells	<i>In vitro</i>	169
Proanthocyanidins	Increasing G1-phase arrest Inhibiting cyclin-dependent kinases (Cdk) Cdk2, Cdk4, Cdk6 and cyclins D1, D2 and E Increasing the protein expression of cyclin-dependent kinase inhibitors (Cdk), Cip1/p21 and Kip1/p27 Enhancing the binding of Cdk-Cdk	In human epidermoid carcinoma A431 cells	<i>In vitro</i>	170

3.14 Effect in the treatment of chronic obstructive pulmonary disease (COPD)

Since a variety of oxidants and free radicals are implicated in the pathogenesis of COPD, it is possible that therapeutic administration of multiple antioxidants will be effective in the treatment of COPD. Various approaches to enhance lung antioxidant capacity and clinical trials of dietary polyphenols in COPD are discussed. Resveratrol, EGCG, and quercetin could inhibit inflammatory gene expression by controlling NF- κ B activation and regulate GSH biosynthesis and chromatin remodel in human airway epithelial A549 cells [171,172]. Curcumin could decrease protein/mRNA expressions of pulmonary type I collagen (Col-I) and TGF- β 1 in rats [173].

3.15 Other bioactive effects

It has been demonstrated that dietary polyphenols have other bioactive effects (Table 11), such as antibacterial activity of Gnemonol B and gnetin E [174], anti-HIV effect of proanthocyanidins [176], hepatoprotective ability of a novel proanthocyanidins IH636 [178], and angiogenesis effect of proanthocyanidins [177].

Table 11. Other bioactive effects of dietary polyphenols.

Type of Activity	Dietary polyphenols	Protective effects and mechanisms	Conditions	Levels	Ref
Antibacterial activity	Gnemonol B and gnetin E	Exhibiting strong antibacterial activities against vancomycin-resistant Enterococci (VRE) and methicillin-resistant Staphylococcus aureus (MRSA)	In Enterococci and Staphylococcus aureus	<i>In vitro</i>	174
	Hydroxytyrosol	Antimycoplasmal activity against <i>M. pneumoniae</i> , <i>M. hominis</i> , and <i>M. fermentans</i>	In <i>Mycoplasma</i>	<i>In vitro</i>	175
Anti-HIV effects	Proanthocyanidins	Downregulating the expression of the HIV-1 entry co-receptors, CCR2b, CCR3 and CCR5	In normal peripheral blood mononuclear cells	<i>In vitro</i>	176
Angiogenesis effect	Proanthocyanidins Resveratrol	Upregulating VEGF expression	In cultured keratinocytes	<i>In vitro</i>	177
Hepato-protective ability	A novel Proanthocyanidins IH636	Increasing the expression of Bcl-xL Attenuating acetaminophen-induced hepatic DNA damage, apoptotic and necrotic cell death of liver cells	In male ICR mice	<i>In vivo</i>	178
	Daidzein	Ameliorating the d-galactosamine-induced increase in malondialdehyde-protein adducts and cytosolic SOD activities	In the rat liver	<i>In vivo</i>	179
	Genistein	Reducing experimental liver damage caused by CCl ₄ by preventing lipid peroxidation and strengthening antioxidant systems	In rats	<i>In vitro</i>	180

4. Prooxidant activity and cellular effects of the phenoxyl radicals of dietary polyphenols

Dietary polyphenols have beneficial antioxidant, anti-inflammatory and anticancer effects. However, at higher doses or under certain conditions these compounds may exert toxic prooxidant activities [181]. Galati *et al.* [182] have observed that dietary polyphenols with phenol rings were metabolized by peroxidase to form prooxidant phenoxyl radicals which, in some cases were sufficiently reactive to cooxidize GSH or NADH accompanied by extensive oxygen uptake and reactive oxygen species formation. Polyphenols with catechol rings also cooxidized ascorbate, likely mediated by semiquinone radicals. Incubation of hepatocytes with dietary polyphenols containing

phenol rings was found to partially oxidize hepatocyte GSH to GSSG while polyphenols with a catechol ring were found to deplete GSH through formation of GSH conjugates.

Dietary polyphenols with phenol rings also oxidized human erythrocyte oxyhemoglobin and caused erythrocyte hemolysis more readily than polyphenols with catechol rings. It is concluded that polyphenols containing a phenol ring are generally more prooxidant than polyphenols containing a catechol ring. Subsequent studies revealed that [183] B-ring catechol-type flavonoids showed swift formation of their two electron oxidized quinone type metabolites, even upon their one electron oxidation by peroxidases. Enzymatic and/or chemical (auto) oxidation of the flavonoid generates the flavonoid semiquinone radical, which may be scavenged by GSH, thereby regenerating the flavonoid and generating the thiyl radical of glutathione. This thiyl radical may react with GSH to generate a disulfide radical anion which rapidly reduces molecular oxygen to superoxide anion radicals.

Huisman *et al.* [184] found that wine polyphenols and ethanol do not significantly scavenge superoxide nor affect endothelial nitric oxide production. Studies showed that flavonoids can induce oxidative damage and nick DNA via the production of radicals in the presence of Cu and O₂. Al, Zn, Ca, Mg and Cd have been found to stimulate phenoxy radical-induced lipid peroxidation [185]. As a result of such enzymatic as well as non-enzymatic antioxidant reactions, phenoxy radicals are formed as the primary oxidized products. Phenoxy radicals can initiate lipid peroxidation. It is concluded that the prooxidant cytotoxicity of diet polyphenols is due to formation of ROS [186], role of phenoxy radical/phenol redox couple [187], and stimulation by metals [185].

5. Bioavailability of dietary polyphenols

Polyphenols are the most abundant antioxidants in the human diet. They show a considerable structural diversity, which largely influences their bioavailability [188]. The biological properties of polyphenols depend on the amount consumed and on their bioavailability. Bioavailability appears to differ greatly between the various polyphenols, and the most abundant polyphenols in our diet are not necessarily those leading to the highest concentrations of active metabolites in target tissues [189]. Both isoflavones and phenolic acids like caffeic acid and gallic acid are the most well absorbed polyphenols, followed by catechins, flavanones, and quercetin glucosides, but with different kinetics. The least well-absorbed polyphenols are large molecular weight polyphenols such as the proanthocyanidins, the galloylated tea catechins, and the anthocyanins [190].

Ellagic acid was detected in human plasma at a maximum concentration (31.9 ng/mL) after 1 h postingestion [191]. Absorption of flavanols such as catechins was enhanced when tea polyphenols were administered as a green tea supplement in capsule form when consumed in the absence of food and led to a small but significant increase in plasma antioxidant activity compared with when tea polyphenols were consumed as black tea or green tea [192,193]. No differences were found in plasma EGCG concentrations and trolox equivalents determined by the trolox equivalent antioxidant capacity assay after administration as a single large dose in the form of either purified EGCG or as green tea extract (Polyphenon E) [194]. Hydroxytyrosol, the major olive oil phenolic compound, is dose-dependently absorbed from olive oil [195]. Tuck *et al.* showed that hydroxytyrosol intravenously and orally administered oil-based dosings resulted in significantly greater elimination of the phenolics in urine within 24 h than the oral, aqueous dosing method. Oral bioavailability estimates of hydroxyl-

tyrosol when administered in an olive oil solution and when dosed as an aqueous solution was 99% and 75%, respectively [13].

Once absorbed, polyphenols are conjugated to glucuronide, sulphate and methyl groups in the gut mucosa and inner tissues. Non-conjugated polyphenols are virtually absent in plasma. Such reactions facilitate their excretion and limit their potential toxicity. EGCG and ECG were present in plasma mostly as the free form, whereas epicatechin and epigallocatechin were mostly present as the glucuronide and sulfate conjugates [192]. Recent data suggest that beta-glucosidases and maybe also lactase phlorizin hydrolase (LPH) in the small intestine are capable of hydrolysing flavonoid glucosides and these compounds are thus taken up as the free aglycon and not as the intact glycosides [196]. It has been reported that around 98% of hydroxytyrosol is present in plasma and urine in conjugated forms, mainly glucuronconjugates, suggesting an extensive first pass intestinal/ hepatic metabolism of the ingested primary forms [197-199] and the 3-*O*-glucuronide of hydroxytyrosol shows stronger activity as a radical scavenger than hydroxytyrosol itself [200]. The major metabolites identified in *in vitro* and *in vivo* studies were an Omethylated derivative of hydroxytyrosol, glucuronides of hydroxytyrosol and a novel glutathionyl conjugate of hydroxytyrosol [200,201]. It has been recently reported that hydroxytyrosol and its metabolites are capable of binding human LDL after olive oil ingestion [202].

The polyphenols reaching the colon are extensively metabolised by the microflora into a wide array of low molecular weight phenolic acids. It has been shown that the plasma concentrations of total metabolites ranged from 0 to 4 $\mu\text{mol/L}$ with an intake of 50 mg aglycone equivalents, and the relative urinary excretion ranged from 0.3% to 43% of the ingested dose, depending on the polyphenol [189]. The biological properties of both conjugated derivatives and microbial metabolites will be essential to better assess the health effects of dietary polyphenols. Alternatively, some health effects of polyphenols may not require their absorption through the gut barrier. Their role as iron chelators in the gut lumen is briefly discussed. Tannic acid and catechin both interact with the gut but only catechin appears able to traverse the gut. In addition, they provide evidence for binding of tannic acid and catechin by endogenous proteins in the intestinal lumen. This may limit their absorption from the small intestine [203].

6. Conclusions

Consumption of polyphenol-rich fruits, vegetables, and beverages derived from plants, such as cocoa, red wine and tea, represents a diet beneficial to human health. Some dietary polyphenols possess antioxidative and anti-inflammatory properties, to some extent, contributing to their cancer chemopreventive potential. These phenolic substances have the ability to abrogate various biochemical processes induced or mediated by the tumor promoters. Some dietary polyphenols also induce apoptosis in premalignant or cancerous cells, and suppress growth and proliferation of various types of tumor cells via induction of apoptosis or arrest of a specific phase of the cell cycle.

However, the specific mechanism(s) by which these compounds affect human health remains unclear, despite extensive research conducted in this area in recent years. Most of that research has focused on the antioxidant properties of dietary polyphenols, which are well characterized and well established *in vitro*. The *in vitro* data often conflict with results obtained from *in vivo* studies on the antioxidant capacity of plasma or the resistance of plasma and lipoproteins to oxidation *ex vivo* after the consumption of polyphenols-rich foods by human subjects. These inconsistencies between the *in*

in vitro and the *in vivo* data are likely explained by the limited bioavailability of dietary polyphenols and their extensive metabolism in humans. Most of them exert multifacet action, and any clinical applications using these substances should be based on the precise understanding of the physiologically relevant action mechanisms.

Acknowledgements

This project was supported by a grant from the National Natural Science Foundation of P.R.China (No. 30472072)

References

1. Gutteridge, J.M. Free radicals in diseases processes: a compilation of cause and consequence. *Free Radic. Res. Commun.* **1993**, *19*, 141-158.
2. Kehrer, J.P. Free radicals as mediators of tissue injury and disease. *Crit. Rev. Toxicol.* **1993**, *23*, 21- 48.
3. Becker, L.B. New concepts in reactive oxygen species and cardiovascular reperfusion physiology. *Cardiovasc. Res.* **2004**, *61*, 461-470.
4. Hayes, J.D.; McLellan, L.I. Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defences against oxidative stress. *Free Radic. Res.* **1999**, *31*, 273-300.
5. Masella, R.; Di Benedetto, R.; Vari, R.; Filesi, C.; Giovannini, C. Novel mechanisms of natural antioxidant compounds in biological systems: involvement of glutathione and glutathione-related enzymes. *J. Nutr. Biochem.* **2005**, *16*, 577-586.
6. Hartman, R.E.; Shah, A.; Fagan, A.M.; Schwetye, K.E.; Parsadonian, M.; Schulman, R. N.; Beth Finn, M.; Holtzman, D.M. Pomegranate juice decreases amyloid load and improves behavior in a mouse model of Alzheimer's disease. *Neurobiol. Dis.* **2006**, *24*, 506-515.
7. Hollman, P.C.; van Trijp, J.M.; Buysman, M.N.; van der Gaag, M.S.; Mengelers, M.J.; de Vries, J.H.; Katan, M.B. Relative bioavailability of the antioxidant flavonoid quercetin from various foods in man. *FEBS Lett.* **1997**, *418*, 152-156.
8. Shen, S.Q.; Zhang, Y.; Xiang, J.J.; Xiong, C.L. Protective effect of curcumin against liver warm ischemia/reperfusion injury in rat model is associated with regulation of heat shock protein and antioxidant enzymes. *World J. Gastroenterol.* **2007**, *13*, 1953-1961.
9. Molina, M.F.; Sanchez-Reus, I.; Iglesias, I.; Benedi, J. Quercetin, a flavonoid antioxidant, prevents and protects against ethanol-induced oxidative stress in mouse liver. *Biol. Pharm. Bull.* **2003**, *26*, 1398-1402.
10. Chen, C.; Yu, R.; Owuor, E.D.; Kong, A.N. Activation of antioxidant response element (ARE), mitogen-activated protein kinases (MAPKs) and caspases by major green tea polyphenol components during cell survival and death. *Arch. Pharm. Res.* **2000**, *23*, 605-612.
11. Butterfield, D.A.; Castegna, A.; Pocernich, C. B.; Drake, J.; Scapagninib, G.; Calabrese, V. Nutritional approaches to combat oxidative stress in Alzheimer's disease. *J. Nutr. Biochem.* **2002**, *13*, 444-461.
12. Schaffer, S.; Podstawa, M.; Visioli, F.; Bogani, P.; Müller, W.E.; Eckert, G.P. Hydroxytyrosol-rich olive mill wastewater extract protects brain cells *in vitro* and *ex vivo*. *J. Agric. Food Chem.* **2007**, *55*, 5043-5049.

13. Tuck, K.L.; Freeman, M.P.; Hayball, P.J.; Stretch, G.L.; Stupans, I. The in vivo fate of hydroxytyrosol and tyrosol, antioxidant phenolic constituents of olive oil, after intravenous and oral dosing of labeled compounds to rats. *J. Nutr.* **2001**, *131*, 1993-1996.
14. Rice-Evans, C.A.; Miller, N.J.; and Paganga, G. Antioxidant properties of phenolic compounds. *Trends Plant Sci.* **1997**, *2*, 152-159.
15. Nielsen, I.L.; Dragsted, L.O.; Ravn-Haren, G.; Freese, R.; Rasmussen, S.E. Absorption and excretion of black currant anthocyanins in humans and watanabe heritable hyperlipidemic rabbits. *J. Agric. Food Chem.* **2003**, *51*, 2813-2820.
16. Bub, A.; Watzl, B.; Heeb, D.; Rechkemmer, G.; Briviba, K. Malvidin-3-glucoside bioavailability in humans after ingestion of red wine, dealcoholized red wine and red grape juice. *Eur. J. Nutr.* **2001**, *40*, 113-120.
17. McAnlis, G.T.; McEneny, J.; Pearce, J.; Young, I.S. Absorption and antioxidant effects of quercetin from onions, in man. *Eur. J. Clin. Nutr.* **1999**, *53*, 92-96.
18. Manach, C.; Morand, C.; Gil-Izquierdo, A.; Bouteloup-Demange, C.; Remesy, C. Bioavailability in humans of the flavanones hesperidin and narirutin after the ingestion of two doses of orange juice. *Eur. J. Clin. Nutr.* **2003**, *57*, 235-242.
19. Lotito, S.B.; Frei, B. Consumption of flavonoid-rich foods and increased plasma antioxidant capacity in humans: Cause, consequence, or epiphenomenon? *Free Radic. Biol. Med.* **2006**, *41*, 1727-1746.
20. Erlund, I.; Meririnne, E.; Alfthan, G.; Aro, A. Plasma kinetics and urinary excretion of the flavanones naringenin and hesperetin in humans after ingestion of orange juice and grapefruit juice. *J. Nutr.* **2001**, *131*, 235-241.
21. Henning, S.M.; Niu, Y.; Liu, Y.; Lee, N.H.; Hara, Y.; Thames, G.D.; Minutti, R.R.; Carpenter, C.L.; Wang, H.; Heber, D. Bioavailability and antioxidant effect of epigallocatechin gallate administered in purified form versus as green tea extract in healthy individuals. *J. Nutr. Biochem.* **2005**, *16*, 610-616.
22. Widlansky, M.E.; Duffy, S.J.; Hamburg, N.M.; Gokce, N.; Warden, B.A.; Wiseman, S.; Keaney, Jr., J.F.; Frei, B.; Vita, J.A. Effects of black tea consumption on plasma catechins and markers of oxidative stress and inflammation in patients with coronary artery disease. *Free Radic. Biol. Med.* **2005**, *38*, 499-506.
23. Bell, J.R.; Donovan, J.L.; Wong, R.; Waterhouse, A.L.; German, J.B.; Walzem, R. L.; Kasim-Karakas, S.E. (+)-Catechin in human plasma after ingestion of a single serving of reconstituted red wine. *Am. J. Clin. Nutr.* **2000**, *71*, 103-108.
24. Holt, R.R.; Lazarus, S.A.; Sullards, M.C.; Zhu, Q.Y.; Schramm, D.D.; Hammerstone, J.F.; Fraga, C.G.; Schmitz, H.H.; Keen, C.L. Procyanidin dimer B2 [epicatechin-(4 β -8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *Am. J. Clin. Nutr.* **2002**, *76*, 798-804.
25. Lotito, S.B.; Frei, B. Consumption of flavonoid-rich foods and increased plasma antioxidant capacity in humans: Cause, consequence, or epiphenomenon? *Free Radic. Biol. Med.* **2006**, *41*, 1727-1746.

26. Schwarz, D. and Roots, I. In vitro assessment of inhibition by natural polyphenols of metabolic activation of procarcinogens by human CYP1A1. *Biochem. Biophys. Res. Commun.* **2003**, *303*, 902-907.
27. Gonthier, M.P.; Remesy, C.; Scalbert, A.; Cheynier, V.; Souquet, J.M.; Poutanen, K.; Aura, A.M. Microbial metabolism of caffeic acid and its esters chlorogenic and caftaric acids by human faecal microbiota in vitro. *Biomed. Pharmacother.* **2006**, *60*, 536-540.
28. Seeram, N. P.; Lee, R.; Heber, D. Bioavailability of ellagic acid in human plasma after consumption of ellagitannins from pomegranate (*Punica granatum* L.) juice. *Clin. Chim. Acta* **2004**, *348*, 63-68.
29. Rangkadilok, N.; Sitthimonchai, S.; Worasuttayangkurn, L.; Mahidol, C.; Ruchirawat, M.; Satayavivad, J. Evaluation of free radical scavenging and antityrosinase activities of standardized longan fruit extract. *Food Chem. Toxicol.* **2007**, *45*, 328-336.
30. Ray, P.S.; Maulik, G.; Cordis, G.A.; Bertelli, A.A.; Bertelli, A.; Das, D.K. The red wine antioxidant resveratrol protects isolated rat hearts from ischemia reperfusion injury. *Free Radic. Biol. Med.* **1999**, *27*, 160-169.
31. Zhang, Y.; Liu, Y.; Wang, T.; Li, B.; Li, H.; Wang, Z.; Yang, B. Resveratrol, a natural ingredient of grape skin: Antiarrhythmic efficacy and ionic mechanisms. *Biochem. Biophys. Res. Commun.* **2006**, *340*, 1192-1199.
32. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y. Tannins and human health: a review. *Crit. Rev. Food Sci. Nutr.* **1998**, *38*, 421-464.
33. Sharma, R.A.; Gescher, A.J.; Steward, W.P. Curcumin: The story so far. *Eur. J. Cancer* **2005**, *41*, 1955-1968.
34. Hong, J.; Smith, T.J.; Ho, C.T.; August, D.A.; Yang, C.S. Effects of purified green and black tea polyphenols on cyclooxygenase- and lipoxygenase-dependent metabolism of arachidonic acid in human colon mucosa and colon tumor tissues. *Biochem. Pharmacol.* **2001**, *62*, 1175-1183.
35. Fki, I.; Sahnoun, Z.; Sayadi, S. Hypocholesterolemic effects of phenolic extracts and purified hydroxytyrosol recovered from olive mill wastewater in rats fed a cholesterol-rich diet. *J. Agric. Food Chem.* **2007**, *55*, 624-631.
36. Kohyama, N.; Nagata, T.; Fujimoto, S.; Sekiya, K. Inhibition of arachidonate lipoxygenase activities by 2-(3, 4-dihydroxyphenyl) ethanol, a phenolic compound from olives. *Biosci. Biotechnol. Biochem.* **1997**, *61*, 347-350.
37. Du, Y.; Guo, H.; Lou, H. Grape seed polyphenols protect cardiac cells from apoptosis via induction of endogenous antioxidant enzymes. *J. Agric. Food Chem.* **2007**, *55*, 1695-1701.
38. Appiah-Opong, R.; Commandeur, J.N.; van Vugt-Lussenburg, B.; Vermeulen, N.P. Inhibition of human recombinant cytochrome P450s by curcumin and curcumin decomposition products. *Toxicology* **2007**, *235*, 83-91.
39. Zheng, J.; Ramirez, V.D. Inhibition of mitochondrial proton F₀F₁-ATPase/ATP synthase by polyphenolic phytochemicals. *Br. J. Pharmacol.* **2000**, *130*, 1115-1123.
40. Nishinaka, T.; Ichijo, Y.; Ito, M.; Kimura, M.; Katsuyama, M.; Iwata, K.; Miura, T.; Terada, T.; Yabe-Nishimura, C. Curcumin activates human glutathione S-transferase P1 expression through antioxidant response element. *Toxicol Lett.* **2007**, *170*, 238-247.

41. Gil, B.; Sanz, M.J.; Terencio, M.C.; Ferrandiz, M.L.; Bustos, G.; Paya M, Gunasegaran, R.; Alcaraz, M.J. Effects of flavonoids on *Naja naja* and human recombinant synovial phospholipase A2 and inflammatory responses in mice. *Life Sci.* **1994**, *54*, 333-338.
42. Huang, J.; de Paulis, T.; May, J.M. Antioxidant effects of dihydrocaffeic acid in human EA.hy926 endothelial cells. *J. Nutr. Biochem.* **2004**, *15*, 722-729.
43. Kerry, N.; Rice-Evans, C. Inhibition of peroxynitrite-mediated oxidation of dopamine by flavonoid and phenolic antioxidants and their structural relationship. *J. Neurochem.* **1999**, *73*, 247-253.
44. Alía, M.; Ramos, S.; Mateos, R.; Granado-Serrano, A.B.; Bravo, L.; Goya, L. Quercetin protects human hepatoma HepG2 against oxidative stress induced by tert-butyl hydroperoxide. *Toxicol. Appl. Pharmacol.* **2006**, *212*, 110-118.
45. Valerio, L.G.; Jr.; Kepa, J.K.; Pickwell, G.V.; Quattrochi, L.C. Induction of human NAD(P)H:quinone oxidoreductase (NQO1) gene expression by the flavonol quercetin. *Toxicol. Lett.* **2001**, *119*, 49-57.
46. Motohashi, H.; Yamamoto, M. Nrf2-Keap1 defines a physiologically important stress response mechanism. *Trends Mol. Med.* **2004**, *10*, 549-557.
47. Scharf, G.; Prustomersky, S.; Knasmuller, S.; Schulte-Hermann, R.; Huber, W.W. Enhancement of glutathione and g-glutamylcysteine synthetase, the rate limiting enzyme of glutathione synthesis, by chemoprotective plant-derived food and beverage components in the human hepatoma cell line HepG2. *Nutr. Cancer* **2003**, *45*, 74-83.
48. Tanigawa, S.; Fujii, M.; Hou, D.X. Action of Nrf2 and Keap1 in ARE-mediated NQO1 expression by quercetin. *Free Radic. Biol. Med.* **2007**, *42*, 1690-1703.
49. Cadenas, S.; Barja, G. Resveratrol, melatonin, vitamin E, and PBN protect against renal oxidative DNA damage induced by the kidney carcinogen KBrO3. *Free Radic. Biol. Med.* **1999**, *26*, 1531-1537.
50. Jang, M.; Pezzuto, J.M. Effects of resveratrol on 12-O-tetradecanoylphorbol -13-acetate -induced oxidative events and gene expression in mouse skin. *Cancer Lett.* **1998**, *134*, 81-89.
51. Dubuisson, J.G.; Dyess, D.L.; Gaubatz, J.W. Resveratrol modulates human mammary epithelial cell O-acetyltransferase, sulfotransferase, and kinase activation of the heterocyclic amine carcinogen N-hydroxy-PhIP. *Cancer Lett.* **2002**, *182*, 27-32.
52. Ciolino, H.P.; Yeh, G.C. Inhibition of aryl hydrocarbon induced cytochrome P-4501A1 enzyme activity and CYP1A1 expression by resveratrol. *Mol. Pharmacol.* **1999**, *56*, 760-767.
53. Casper, R.F.; Quesne, M.; Rogers, I.M.; Shirota, T.; Jolivet, A.; Milgrom, E.; Savouret, J.F. Resveratrol has antagonist activity on the aryl hydrocarbon receptor: implications for prevention of dioxin toxicity. *Mol. Pharmacol.* **1999**, *56*, 784-790.
54. Schewe, T.; Sadik, C.; Klotz, L.O.; Yoshimoto, T.; Kuhn, H.; Sies, H. Polyphenols of cocoa: inhibition of mammalian 15-lipoxygenase. *Biol. Chem.* **2001**, *382*, 1687-1696.
55. Subbaramaiah, K.; Chung, W.J.; Michaluart, P.; Telang, N.; Tanabe, T.; Inoue, H.; Jang, M.; Pezzuto, J.M.; Dannenberg, A.J. Resveratrol inhibits cyclooxygenase-2 transcription and activity in phorbol ester-treated human mammary epithelial cells. *J. Biol. Chem.* **1998**, *273*, 21875-21882.
56. Li, Y.T.; Shen, F.; Liu, B.H.; Cheng, G.F. Resveratrol inhibits matrix metalloproteinase-9 transcription in U937 cells. *Acta Pharmacol. Sin.* **2003**, *24*, 1167-1171.

57. Kaga, S.; Zhan, L.; Matsumoto, M.; Maulik, N. Resveratrol enhances neovascularization in the infarcted rat myocardium through the induction of thioredoxin-1, heme oxygenase-1 and vascular endothelial growth factor. *J. Mol. Cell. Cardiol.* **2005**, *39*, 813-822.
58. Cullen, J.P.; Morrow, D.; Jin, Y.; von Offenbergs Sweeney, N.; Sitzmann, J.V.; Cahill, P.A.; Redmond, E.M. Resveratrol inhibits expression and binding activity of the monocyte chemotactic protein-1 receptor, CCR2, on THP-1 monocytes. *Atherosclerosis* **2007**, (in press)
59. Wang, S.; Wang, X.; Yan, J.; Xie, X.; Fan, F.; Zhou, X.; Han, L.; Chen, J. Resveratrol inhibits proliferation of cultured rat cardiac fibroblasts: Correlated with NO-cGMP signaling pathway. *Eur. J. Pharmacol.* **2007**, *567*, 26-35.
60. Steffen, Y.; Wiswedel, I.; Peter, D.; Schewe, T.; Sies, H. Cytotoxicity of myeloperoxidase/nitrite-oxidized low-density lipoprotein toward endothelial cells is due to a high 7 β -hydroxycholesterol to 7-ketocholesterol ratio. *Free Radic. Biol. Med.* **2006**, *41*, 1139-1150.
61. Petroni, A.; Blasevich, M.; Salami, M.; Papini, N.; Montedoro, G.F.; Galli, C. Inhibition of platelet aggregation and eicosanoid production by phenolic components of olive oil. *Thromb. Res.* **1995**, *78*, 151-160.
62. Léger, C.L.; Carbonneau, M.A.; Michel, F.; Mas, E.; Monnier, L.; Cristol, J.P.; Descomps, B. A thromboxane effect of a hydroxytyrosol-rich olive oil wastewater extract in patients with uncomplicated type I diabetes. *Eur. J. Clin. Nutr.* **2005**, *59*, 727-730.
63. de La Puerta, R.; Ruiz-Gutierrez, V.; Houtl, J.R. Inhibition of leukocyte 5-lipoxygenase by phenolics from virgin olive oil. *Biochem. Pharmacol.* **1999**, *57*, 445-449.
64. Carluccio, M.A.; Siculella, L.; Ancora, M.A.; Massaro, M.; Scoditti, E.; Storelli, C.; Visioli, F.; Distanti, A.; De Caterina, R. Olive oil and red wine antioxidant polyphenols inhibit endothelial activation: antiatherogenic properties of Mediterranean diet phytochemicals. *Arterioscler. Thromb. Vasc. Biol.* **2003**, *23*, 622-629.
65. Manna, C.; Migliardi, V.; Golino, P.; Scognamiglio, A.; Galletti, P.; Chiariello, M.; Zappia, V. Oleuropein prevents oxidative myocardial injury induced by ischemia and reperfusion. *J. Nutr. Biochem.* **2004**, *15*, 461-466.
66. Gouedard, C.; Barouki, R.; Morel, Y. Dietary polyphenols increase paraoxonase 1 gene expression by an aryl hydrocarbon receptor-dependent mechanism. *Mol. Cell Biol.* **2004**, *24*, 5209-5222.
67. Nair, M.P.; Kandaswami, C.; Mahajan, S.; Chadha, K.C.; Chawda, R.; Nair, H.; Kumar, N.; Nair, R.E.; Schwartz, S.A. The flavonoid, quercetin, differentially regulates Th-1 (IFN gamma) and Th-2 (IL4) cytokine gene expression by normal peripheral blood mononuclear cells. *Biochim. Biophys. Acta* **2002**, *1593*, 29-36.
68. Myhrstad, M.C.; Carlsen, H.; Nordstrom, O.; Blomhoff, R.; Moskaug, J.O. Flavonoids increase the intracellular glutathione level by transactivation of the gamma-glutamylcysteine synthetase catalytical subunit promoter. *Free Radic. Biol. Med.* **2002**, *32*, 386-393.
69. Lo, H.M.; Hung, C.F.; Huang, Y.Y.; Wu, W.B. Tea polyphenols inhibit rat vascular smooth muscle cell adhesion and migration on collagen and laminin via interference with cell-ECM interaction. *J. Biomed. Sci.* **2007**, (in press)
70. Mizushige, T.; Mizushige, K.; Miyatake, A.; Kishida, T.; Ebihara, K. Inhibitory effects of soy isoflavones on cardiovascular collagen accumulation in rats. *J. Nutr. Sci. Vitaminol. (Tokyo)* **2007**, *53*, 48-52.

71. Tikkanen, M.J.; Adlercreutz, H. Dietary soy-derived isoflavone phytoestrogens. Could they have a role in coronary heart disease prevention? *Biochem.Pharmacol.* **2000**, *60*, 1-5.
72. Schramm, D.D.; Wang, J.F.; Holt, R.R.; Ensunsa, J.L.; Gonsalves, J.L.; Lazarus, S.A.; Schmitz, H.H.; German, J.B.; Keen, C.L. Chocolate procyanidins decrease the leukotriene-prostacyclin ratio in humans and human aortic endothelial cells. *Am. J. Clin. Nutr.* **2001**, *73*, 36-40.
73. Dedoussis, G.V.; Kaliora, A.C.; Psarras, S.; Chiou, A.; Mylona, A.; Papadopoulos, N.G.; Andrikopoulos, N.K. Antiatherogenic effect of Pistacia lentiscus via GSH restoration and downregulation of CD36 mRNA expression. *Atherosclerosis* **2004**, *174*, 293-303.
74. Sato, M.; Bagchi, D.; Tosaki, A.; Das, D.K. Grape seed proanthocyanidin reduces cardiomyocyte apoptosis by inhibiting ischemia-reperfusion-induced activation of JNK-1 and c-JUN. *Free Radic. Biol. Med.* **2001**, *31*, 729-737.
75. Dasgupta, B.; Milbrandt, J. Resveratrol stimulates AMP kinase activity in neurons. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 7217-7222.
76. Chavez, E.; Reyes-Gordillo, K.; Segovia, J.; Shibayama, M.; Tsutsumi, V.; Vergara, P.; Moreno, M.G.; Muriel, P. Resveratrol prevents fibrosis, NF-kappaB activation and TGF-beta increases induced by chronic CCl (4) treatment in rats. *J. Appl. Toxicol.* **2007**, (in press)
77. Bastianetto, S.; Brouillette, J.; Quirion, R. Neuroprotective effects of natural products: interaction with intracellular kinases, amyloid peptides and a possible role for transthyretin. *Neurochem. Res.* **2007**, (in press)
78. Okawara, M.; Katsuki, H.; Kurimoto, E.; Shibata, H.; Kume, T.; Akaike, A. Resveratrol protects dopaminergic neurons in midbrain slice culture from multiple insults. *Biochem. Pharmacol.* **2007**, *73*, 550-560.
79. Kim, S.J.; Jeong, H.J.; Lee, K.M.; Myung, N.Y.; An, N.H.; Mo Yang, W.; Kyu Park, S.; Lee, H.J.; Hong, S.H.; Kim, H.M.; Um, J.Y. Epigallocatechin-3-gallate suppresses NF-kappaB activation and phosphorylation of p38 MAPK and JNK in human astrocytoma U373MG cells. *J. Nutr. Biochem.* **2007**, (in press)
80. de Boer, V.C.; de Goffau, M.C.; Arts, I.C.; Hollman, P.C.; Keijer, J. SIRT1 stimulation by polyphenols is affected by their stability and metabolism. *Mech. Ageing Dev.* **2006**, *127*, 618-627.
81. Levites, Y.; Amit, T.; Youdim, M.B.; Mandel, S. Involvement of protein kinase C activation and cell survival/cell cycle genes in green tea polyphenol (-)-epigallocatechin-3-gallate neuron-protective action. *J. Biol. Chem.* **2002**, *277*, 30574-30580.
82. Mercer, L. D.; Kelly, B.L.; Horne, M. K.; Beart, P.M. Dietary polyphenols protect dopamine neurons from oxidative insults and apoptosis: investigations in primary rat mesencephalic cultures. *Biochem. Pharmacol.* **2005**, *69*, 339-345.
83. Schroeter, H.; Spencer, J.P.; Rice-Evans, C.; Williams, R.J. Flavonoids protect neurons from oxidized low-density lipoprotein-induced apoptosis involving c-Jun N-terminal kinase (JNK), c-jun and caspase-3. *Biochem. J.* **2001**, *358*, 547-557.
84. Garcia-Alloza, M.; Borrelli, L.A.; Rozkalne, A.; Hyman, B.T.; Bacskai, B.J. Curcumin labels amyloid pathology in vivo, disrupts existing plaques, and partially restores distorted neurites in an Alzheimer mouse model. *J. Neurochem.* **2007**, *102*, 1095-1104.

85. Mao, T.K.; Powell, J.; Van de Water, J.; Keen, C.L.; Schmitz, H.H.; Hammerstone, J.F.; Eric Gershwin, M. The effect of cocoa procyanidins on the transcription and secretion of interleukin β in peripheral blood mononuclear cells. *Life Sci.* **2000**, *66*, 1377-1386.
86. Kawai, K.; Tsuno, N.H.; Kitayama, J.; Okaji, Y.; Yazawa, K.; Asakage, M.; Sasaki, S.; Watanabe, T.; Takahashi, K.; Nagawa, H. Epigallocatechin gallate induces apoptosis of monocytes. *J. Allergy Clin. Immunol.* **2005**, *115*, 186-191.
87. Kawai, K.; Tsuno, N.H.; Kitayama, J.; Okaji, Y.; Yazawa, K.; Asakage, M.; Hori, N.; Watanabe, T.; Takahashi, K.; Nagawa, H. Epigallocatechin gallate attenuates adhesion and migration of CD8⁺ T cells by binding to CD11b. *J. Allergy Clin. Immunol.* **2004**, *113*, 1211-1217.
88. Shakibaei, M.; John, T.; Seifarth, C.; Mobasheri, A. Resveratrol inhibits IL-1 β -induced stimulation of caspase-3 and cleavage of PARP in human articular chondrocytes in vitro. *Ann. N. Y. Acad. Sci.* **2007**, *1095*, 554-563.
89. Tsai, S.H.; Lin-Shiau, S.Y.; Lin, J.K. Suppression of nitric oxide synthase and the down-regulation of the activation of NF- κ B in macrophages by resveratrol. *Br. J. Pharmacol.* **1999**, *126*, 673-680.
90. Nonn, L.; Duong, D.; Peehl, D.M. Chemopreventive anti-inflammatory activities of curcumin and other phytochemicals mediated by MAP kinase phosphatase-5 in prostate cells. *Carcinogenesis* **2007**, *28*, 1188-1196.
91. Gerritsen, M.E.; Carley, W.W.; Ranges, G.E.; Shen, C.P.; Phan, S.A.; Ligon, G.F.; Perry, C.A. Flavonoids inhibit cytokine-induced endothelial cell adhesion protein gene expression. *Am. J. Pathol.* **1995**, *147*, 278-292.
92. Choi, J.S.; Choi, Y.J.; Park, S.H.; Kang, J.S.; Kang, Y.H. Flavones mitigate tumor necrosis factor- α -induced adhesion molecule upregulation in cultured human endothelial cells: role of nuclear factor- κ B. *J. Nutr.* **2004**, *134*, 1013-1019.
93. van Meeteren, M.E.; Hendriks, J.J.; Dijkstra, C.D.; van Tol, E.A. Dietary compounds prevent oxidative damage and nitric oxide production by cells involved in demyelinating disease. *Biochem. Pharmacol.* **2004**, *67*, 967-975.
94. Youdim, K.A.; McDonald, J.; Kalt, W.; Joseph, J.A. Potential role of dietary flavonoids in reducing microvascular endothelium vulnerability to oxidative and inflammatory insults. *J. Nutr. Biochem.* **2002**, *13*, 282-288.
95. Camacho-Barquero, L.; Villegas, I.; Sanchez-Calvo, J.M.; Talero, E.; Sanchez-Fidalgo, S.; Motilva, V.; Alarcon de la Lastra, C. Curcumin, a Curcuma longa constituent, acts on MAPK p38 pathway modulating COX-2 and iNOS expression in chronic experimental colitis. *Int. Immunopharmacol.* **2007**, *7*, 333-342.
96. Rahman I, Biswas SK, Kirkham PA. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem. Pharmacol.* **2006**, *72*, 1439-1452.
97. Kim, H.Y.; Park, E.J.; Joe, E.H.; Jou, I. Curcumin suppresses Janus kinase-STAT inflammatory signaling through activation of src homology 2 domain-containing tyrosine phosphatase 2 in brain microglia. *J. Immunol.* **2003**, *171*, 6072-6079.
98. Fabiani, R.; De Bartolomeo, A.; Rosignoli, P.; Servili, M.; Montedoro, G.F.; Morozzi, G. Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through G1 cell cycle arrest and apoptosis. *Eur. J. Cancer Prev.* **2002**, *11*, 351-358.

99. Fuggetta, M.P.; Lanzilli, G.; Tricarico, M.; Cottarelli, A.; Falchetti, R.; Ravagnan, G.; Bonmassar, E. Effect of resveratrol on proliferation and telomerase activity of human colon cancer cells in vitro. *J. Exp. Clin. Cancer Res.* **2006**, *25*, 189-193.
100. Kuo, P.L.; Chiang, L.C.; Lin, C.C. Resveratrol-induced apoptosis is mediated by p53-dependent pathway in HepG2 cells. *Life Sci.* **2002**, *72*, 23-34.
101. Pozo-Guisado, E.; Lorenzo-Benayas, M.J.; Fernandez-Salguero, P.M. Resveratrol modulates the phosphoinositide 3-kinase pathway through an estrogen receptor alpha-dependent mechanism: relevance in cell proliferation. *Int. J. Cancer* **2004**, *109*, 167-173.
102. Kundu, J.K.; Chun, K.S.; Kim, S.O.; Surh, Y.J. Resveratrol inhibits phorbol ester-induced cyclooxygenase-2 expression in mouse skin: MAPKs and AP-1 as potential molecular targets. *Biofactors* **2004**, *21*, 33-39.
103. Yoon, S.H.; Kim, Y.S.; Ghim, S.Y.; Song, B.H.; Bae, Y.S. Inhibition of protein kinase CKII activity by resveratrol, a natural compound in red wine and grapes. *Life Sci.* **2002**, *71*, 2145-2152.
104. Slater, S.J.; Seiz, J.L.; Cook, A.C.; Stagliano, B.A.; Buzas, C.J. Inhibition of protein kinase C by resveratrol. *Biochim. Biophys. Acta* **2003**, *1637*, 59-69.
105. Li, H.; Cheng, Y.; Wang, H.; Sun, H.; Liu, Y.; Liu, K.; Peng, S. Inhibition of nitrobenzene-induced DNA and hemoglobin adductions by dietary constituents. *Appl. Radiat. Isot.* **2003**, *58*, 291-298.
106. Grace, S.C.; Salgo, M.G.; Pryor, W.A. Scavenging of peroxynitrite by a phenolic/peroxidase system prevents oxidative damage to DNA. *FEBS Lett.* **1998**, *426*, 24-28.
107. Lee, L.T.; Huang, Y.T.; Hwang, J.J.; Lee, P.P.; Ke, F.C.; Nair, M.P.; Kanadaswam, C.; Lee, M.T. Blockade of the epidermal growth factor receptor tyrosine kinase activity by quercetin and luteolin leads to growth inhibition and apoptosis of pancreatic tumor cells. *Anticancer Res.* **2002**, *22*, 1615-1627.
108. Cooray, H.C.; Janvilisri, T.; van Veen, H.W.; Hladky, S.B. and Barrant, M.A. Interaction of the breast cancer resistance protein with plant polyphenols. *Biochem. Biophys. Res. Commun.* **2004**, *317*, 269-275.
109. Kunnumakkara, A.B.; Guha, S.; Krishnan, S.; Diagaradjane, P.; Gelovani, J.; Aggarwal, B.B. Curcumin potentiates antitumor activity of gemcitabine in an orthotopic model of pancreatic cancer through suppression of proliferation, angiogenesis, and inhibition of nuclear factor-kappaB-regulated gene products. *Cancer Res.* **2007**, *67*, 3853-3861.
110. Collett, G.P.; Campbell, F.C. Curcumin induces c-jun N-terminal kinase-dependent apoptosis in HCT116 human colon cancer cells. *Carcinogenesis* **2004**, *25*, 2183-2189.
111. Anto, R.J.; Mukhopadhyay, A.; Denning, K.; Aggarwal, B.B. Curcumin (diferuloylmethane) induces apoptosis through activation of caspase-8, BID cleavage and cytochrome c release: its suppression by ectopic expression of Bcl-2 and Bcl-xL. *Carcinogenesis* **2002**, *23*, 143-150.
112. Aoki, H.; Takada, Y.; Kondo, S.; Sawaya, R.; Aggarwal, B.; Kondo, Y. Aoki, H.; Takada, Y.; Kondo, S.; Sawaya, R.; Aggarwal, B.; Kondo, Y. Evidence that curcumin suppresses the growth of malignant gliomas in vitro and in vivo through induction of autophagy: role of Akt and ERK signaling pathways. *Mol. Pharmacol.* **2007**, *72*, 29-39.

113. Nardini, M.; Leonardi, F.; Scaccini, C.; Virgili, F. Modulation of ceramide-induced NF- κ B binding activity and apoptotic response by caffeic acid in U937 cells: comparison with other antioxidants. *Free Radic. Biol. Med.* **2001**, *30*, 722-733.
114. Naasani, I.; Oh-Hashi, F.; Oh-Hara, T.; Feng, W.Y.; Johnston, J.; Chan, K.; Tsuruo, T. Blocking telomerase by dietary polyphenols is a major mechanisms for limiting the growth of human cancer cells in vitro and in vivo. *Cancer Res.* **2003**, *63*, 824-830.
115. Jung, J.Y.; Mo, H.C.; Yang, K.H.; Jeong, Y.J.; Yoo, H.G.; Choi, N.K.; Oh, W.M.; Oh, H.K.; Kim, S.H.; Lee, J.H.; Kim, H.J.; Kim, W.J. Inhibition by epigallocatechin gallate of CoCl₂-induced apoptosis in rat PC12 cells. *Life Sci.* **2007**, *80*, 1355-1363.
116. Khanduja, K.L.; Avti, P.K.; Kumar, S.; Mittal, N.; Sohi, K.K.; Pathak, C.M. Anti-apoptotic activity of caffeic acid, ellagic acid and ferulic acid in normal human peripheral blood mononuclear cells: A Bcl-2 independent mechanism. *Biochim. Biophys. Acta* **2006**, *1760*, 283-289.
117. Choi, Y.J.; Jeong, Y.J.; Lee, Y.J.; Kwon, H.M.; Kang, Y.H. (-) Epigallocatechin gallate and quercetin enhance survival signaling in response to oxidant-induced human endothelial apoptosis. *J. Nutr.* **2005**, *135*, 707-713.
118. Bharti, A.C.; Donato, N.; Aggarwal, B.B. Curcumin (diferuloylmethane) inhibits constitutive and IL-6-inducible STAT3 phosphorylation in human multiple myeloma cells. *J. Immunol.* **2003**, *171*, 3863-3871.
119. Yuste, P.; Longstaff, M.; McCorquodale, C. The effect of proanthocyanidin-rich hulls and proanthocyanidin extracts from bean (*Vicia faba* L.) hulls on nutrient digestibility and digestive enzyme activities in young chicks. *Br. J. Nutr.* **1992**, *67*, 57-65.
120. Longstaff, M.; McNab, J.M. The inhibitory effects of hull polysaccharides and tannins of field beans (*Vicia faba* L.) on the digestion of amino acids, starch and lipid and on digestive enzyme activities in young chicks. *Br. J. Nutr.* **1991**, *65*, 199-216.
121. Sbarra, V.; Ristorcelli, E.; Petit-Thevenin, J.L.; Teissedre, P.L.; Lombardo, D.; Verine, A. In vitro polyphenol effects on activity, expression and secretion of pancreatic bile salt-dependent lipase. *Biochim. Biophys. Acta* **2005**, *1736*, 67-76.
122. Aggarwal, B.B.; Shisodia, S. Suppression of the nuclear factor- κ B activation pathway by spice-derived phytochemicals: reasoning for seasoning. *Ann. N.Y. Acad. Sci.* **2004**, *1030*, 434-441.
123. Hanamura, T.; Hagiwara, T.; Kawagishi, H. Structural and functional characterization of polyphenols isolated from acerola (*Malpighia emarginata* DC.) fruit. *Biosci. Biotechnol. Biochem.* **2005**, *69*, 280-286.
124. Youn, H.S.; Saitoh, S.I.; Miyake, K.; Hwang, D.H. Inhibition of homodimerization of Toll-like receptor 4 by curcumin. *Biochem. Pharmacol.* **2006**, *72*, 62-69.
125. Pan, M.H.; Lin-Shiau, S.Y.; Lin, J.K. Comparative studies on the suppression of nitric oxide synthase by curcumin and its hydrogenated metabolites through down-regulation of IkappaB kinase and NFkappaB activation in macrophages. *Biochem. Pharmacol.* **2000**, *60*, 1665-1676.
126. Kang, G.; Kong, P.J.; Yuh, Y.J.; Lim, S.Y.; Yim, S.V.; Chun, W.; Kim, S.S. Curcumin suppresses lipopolysaccharide-induced cyclooxygenase-2 expression by inhibiting activator

- protein 1 and nuclear factor kappaB bindings in BV2 microglial cells. *J. Pharmacol. Sci.* **2004**, *94*, 325-328.
127. Kluth, D.; Banning, A.; Paur, I.; Blomhoff, R.; Brigelius-Flohe, R. Modulation of pregnane X receptor-and electrophile responsive element-mediated gene expression by dietary polyphenolic compounds. *Free Radic. Biol. Med.* **2007**, *42*, 315-325.
128. Meeran, S.M.; Katiyar, S.K. Grape seed proanthocyanidins promote apoptosis in human epidermoid carcinoma A431 cells through alterations in Cdk1-Cdk-cyclin cascade, and caspase-3 activation via loss of mitochondrial membrane potential. *Exp. Dermatol.* **2007**, *16*, 405-415.
129. Sharma, S.D.; Meeran, S.M.; Katiyar, S.K. Dietary grape seed proanthocyanidins inhibit UVB-induced oxidative stress and activation of mitogen-activated protein kinases and nuclear factor-kappaB signaling in in vivo SKH-1 hairless mice. *Mol. Cancer Ther.* **2007**, *6*, 995-1005.
130. Spencer, J.P.; Rice-Evans, C.; Williams, R.J. Modulation of pro-survival Akt/PKB and ERK1/2 signalling cascades by quercetin and its in vivo metabolites underlie their action on neuronal viability. *J. Biol. Chem.* **2003**, *278*, 34783-34793.
131. Joy, S.; Siow, R.C.; Rowlands, D.J.; Becker, M.; Wyatt, A.W.; Aaronson, P.I.; Coen, C.W.; Kallo, I.; Jacob, R.; Mann, G.E. The isoflavone Equol mediates rapid vascular relaxation: Ca²⁺-independent activation of endothelial nitric-oxide synthase/Hsp90 involving ERK1/2 and Akt phosphorylation in human endothelial cells. *J. Biol. Chem.* **2006**, *281*, 27335-27345.
132. Chen, C.Y.; Jang, J.H.; Li, M.H.; Surh, Y.J. Resveratrol upregulates heme oxygenase-1 expression via activation of NF-E2-related factor 2 in PC12 cells. *Biochem. Biophys. Res. Commun.* **2005**, *331*, 993-1000.
133. Oak, M.H.; Chataigneau, M.; Keravis, T.; Chataigneau, T.; Beretz, A.; Andriantsitohaina, R.; Stoclet, J.C.; Chang, S.J.; Schini-Kerth, V.B. Red wine polyphenolic compounds inhibit vascular endothelial growth factor expression in vascular smooth muscle cells by preventing the activation of the p38 mitogen-activated protein kinase pathway. *Arterioscler. Thromb. Vasc. Biol.* **2003**, *23*, 1001-1007.
134. Oak, M.H.; El Bedoui, J.; Anglard, P.; Schini-Kerth, V.B. Red wine polyphenolic compounds strongly inhibit pro-matrix metalloproteinase-2 expression and its activation in response to thrombin via direct inhibition of membrane type 1-matrix metalloproteinase in vascular smooth muscle cells. *Circulation* **2004**, *110*, 1861-1867
135. Iijima, K.; Yoshizumi, M.; Hashimoto, M.; Akishita, M.; Kozaki, K.; Ako, J.; Watanabe, T.; Ohike, Y.; Son, B.; Yu, J.; Nakahara, K.; Ouchi, Y. Red wine polyphenols inhibit vascular smooth muscle cell migration through two distinct signaling pathways. *Circulation* **2002**, *105*, 2404-2410.
136. Ndiaye, M.; Chataigneau, T.; Chataigneau, M.; Schini-Kerth, V.B. Red wine polyphenols induce EDHF-mediated relaxations in porcine coronary arteries through the redox-sensitive activation of the PI3-kinase/Akt pathway. *Br. J. Pharmacol.* **2004**, *142*, 1131-1136.
137. Martin, S.; Andriambelolon, E.; Takeda, K.; Andriantsitohaina, R. Red wine polyphenols increase calcium in bovine aortic endothelial cells: a basis to elucidate signalling pathways leading to nitric oxide production. *Br. J. Pharmacol.* **2002**, *135*, 1579-1587.
138. Jiménez, R.; López-Sepúlveda, R.; Kadmiri, M.; Romero, M.; Vera, R.; Sánchez, M.; Vargas, F.; O'valle, F.; Zarzuelo, A.; Dueñas, M.; Santos-Buelga, C.; Duarte, J. Polyphenols restore

- endothelial function in DOCA-salt hypertension: Role of endothelin-1 and NADPH oxidase. *Free Radic. Biol. Med.* **2007**, *43*, 462-473.
139. Khan, N.Q.; Lees, D.M.; Douthwaite, J.A.; Carrier, M.J.; Corder, R. Comparison of red wine extract and polyphenol constituents on endothelin-1 synthesis by cultured endothelial cells. *Clin. Sci. (Lond)*. **2002**, *103 Suppl 48*, 72S-75S.
140. Wollny, T.; Chabielska, E.; Malinowska-Zaprzałka, M.; Nazarko, J.; Rozmysłowicz-Szermińska, W.; Buczko, W. Effects of Bulgarian red and white wines on primary hemostasis and experimental thrombosis in rats. *Pol. J. Pharmacol.* **2003**, *55*, 1089-1096.
141. Dell'Agli, M.; Busciala, A.; Bosisio, E. Vascular effects of wine polyphenols. *Cardiovasc. Res.* **2004**, *63*, 593-602.
142. Xu, J.W.; Ikeda, K.; Yamori, Y. Cyanidin-3-glucoside regulates phosphorylation of endothelial nitric oxide synthase. *FEBS Lett.* **2004**, *574*, 176-180.
143. Kim, J.A.; Formoso, G.; Li, Y.; Potenza, M.A.; Marasciulo, F.L.; Montagnani, M.; Quon, M.J. Epigallocatechin gallate, a green tea polyphenol, mediates NO-dependent vasodilation using signaling pathways in vascular endothelium requiring reactive oxygen species and Fyn. *J. Biol. Chem.* **2007**, *282*, 13736-13745.
144. Lorenz, M.; Wessler, S.; Follmann, E.; Michaelis, W.; Düsterhöft, T.; Baumann, G.; Stangl, K.; Stangl, V. A constituent of green tea, epigallocatechin-3-gallate, activates endothelial nitric oxide synthase by a phosphatidylinositol-3-OH-kinase-, cAMP-dependent protein kinase-, and Akt-dependent pathway and leads to endothelial-dependent vasorelaxation. *J. Biol. Chem.* **2004**, *279*, 6190-6195.
145. Maiti, T.K.; Chatterjee, J. and Dasgupta, S. Effect of green tea polyphenols on angiogenesis induced by an angiogenin-like protein. *Biochem. Biophys. Res. Commun.* **2003**, *308*, 64-67.
146. Kalin, R.; Righi, A.; Del Rosso, A.; Bagchi, D.; Generini, S.; Cerinic, M.M.; Das, D.K. Activin, a grape seed-derived proanthocyanidin extract, reduces plasma levels of oxidative stress and adhesion molecules (ICAM-1, VCAM-1 and E-selectin) in systemic sclerosis. *Free Radic. Res.* **2002**, *36*, 819-825.
147. Sen, C.K.; Bagchi, D. Regulation of inducible adhesion molecule expression in human endothelial cells by grape seed proanthocyanidin extract. *Mol. Cell Biochem.* **2001**, *216*, 1-7.
148. Actis-Goretta, L.; Ottaviani, J.I.; Keen, C.L.; Fraga, C.G. Inhibition of angiotensin converting enzyme (ACE) activity by flavan-3-ols and procyanidins. *FEBS Lett.* **2003**, *555*, 597-600.
149. Alvarez, P.; Alvarado, C.; Puerto, M.; Schlumberger, A.; Jiménez, L.; De la Fuente, M. Improvement of leukocyte functions in prematurely aging mice after five weeks of diet supplementation with polyphenol-rich cereals. *Nutrition* **2006**, *22*, 913-921.
150. Bhattacharyya, S.; Mandal, D.; Saha, B.; Sen, G.S.; Das, T.; Sa, G. Curcumin prevents tumor-induced T cell apoptosis through Stat-5a-mediated Bcl-2 induction. *J. Biol. Chem.* **2007**, *282*, 15954-15964.
151. Akiyama, H.; Sato, Y.; Watanabe, T.; Nagaoka, M.H.; Yoshioka, Y.; Shoji, T.; Kanda, T.; Yamada, K.; Totsuka, M.; Teshima, R.; Sawada, J.; Goda, Y.; Maitani, T. Dietary unripe apple polyphenol inhibits the development of food allergies in murine models. *FEBS Lett.* **2005**, *579*, 4485-4491.

152. Kanda, T.; Akiyama, H.; Yanagida, A.; Tanabe, M.; Goda, Y.; Toyoda, M.; Teshima, R.; Saito, Y. Inhibitory effects of apple polyphenol on induced histamine release from RBL-2H3 cells and rat mast cells. *Biosci. Biotechnol. Biochem.* **1998**, *62*, 1284-1289.
153. Kowluru, R.A.; Kanwar, M. Effects of curcumin on retinal oxidative stress and inflammation in diabetes. *Nutr Metab (London)* **2007**, *4*, 8.
154. Johnston, K.; Sharp, P.; Clifford, M.; Morgan, L. Dietary polyphenols decrease glucose uptake by human intestinal Caco-2 cells. *FEBS Lett.* **2005**, *579*, 1653-1657.
155. Kobayashi, Y.; Suzuki, M.; Hideo, S.; Arai, S.; Yukihiro, H.; Suzuki, K.; Miyamoto, Y.; Shimizu, M. Green tea polyphenols inhibit the sodium-dependent glucose transporter of intestinal epithelial cells by a competitive mechanism. *J. Agric. Food Chem.* **2000**, *48*, 5618-5623.
156. Song, J.; Kwon, O.; Chen, S.; Daruwala, R.; Eck, P.; Park, J.B.; Levine, M. Flavonoid inhibition of SVCT1 and GLUT2, intestinal transporters for vitamin C and glucose. *J. Biol. Chem.* **2002**, *277*, 15252-15260.
157. Yoshikawa, M.; Nishida, N.; Shimoda, H.; Takada, M.; Kawahara, Y.; Matsuda, H. Polyphenol constituents from *Salacia* species: quantitative analysis of mangiferin with alpha-glucosidase and aldose reductase inhibitory activities. *Yakugaku Zasshi* **2001**, *121*, 371-378.
158. McDougall GJ, Shpiro F, Dobson P, Smith P, Blake A, Stewart D. Different polyphenolic components of soft fruits inhibits alpha-amylase and alpha-glucosidase. *J. Agric. Food Chem.* **2005**, *53*, 2760-2766.
159. Bhat, K.P.; Lantvit, D.; Christov, K.; Mehta, R.G.; Moon, R.C.; Pezzuto, J.M. Estrogenic and antiestrogenic properties of resveratrol in mammary tumor models. *Cancer Res.* **2001**, *61*, 7456-7463.
160. Bhat, K.P.; Pezzuto, J.M. Resveratrol exhibits cytostatic and antiestrogenic properties with human endometrial adenocarcinoma (Ishikawa) cells. *Cancer Res.* **2001**, *61*, 6137-6144.
161. Otake, Y.; Nolan, A.L.; Walle, U.K.; Walle, T. Quercetin and resveratrol potently reduce estrogen sulfotransferase activity in normal human mammary epithelial cells. *J. Steroid Biochem. Mol. Biol.* **2000**, *73*, 265-270.
162. Reidenberg, M.M. Environmental inhibition of 11b-hydroxysteroid dehydrogenase. *Toxicology* **2000**, *144*, 107-111.
163. Kuo, P.L.; Chiang, L.C.; Lin, C.C. Resveratrol-induced apoptosis is mediated by p53-dependent pathway in HepG2 cells. *Life Sci.* **2002**, *72*, 23-34.
164. Ahmad, N.; Adhami, V. M.; Afaq, F.; Feyes, D. K.; Mukhtar, H. Resveratrol causes WAF-1/p21-mediated G (1)-phase arrest of cell cycle and induction of apoptosis in human epidermoid carcinoma A431 cells. *Clin. Cancer Res.* **2001**, *7*, 1466-1473.
165. Wolter, F.; Akoglu, B.; Clausnitzer, A.; Stein, J. Downregulation of the cyclin D1/Cdk4 complex occurs during resveratrol-induced cell cycle arrest in colon cancer cell lines. *J. Nutr.* **2001**, *131*, 2197-2203.
166. Adhami, V.M.; Afaq, F.; Ahmad, N. Involvement of the retinoblastoma (pRb)-E2F/DP pathway during antiproliferative effects of resveratrol in human epidermoid carcinoma (A431) cells. *Biochem. Biophys. Res. Commun.* **2001**, *288*, 579-585.

167. Joe, A.K.; Liu, H.; Suzui, M.; Vural, M. E.; Xiao, D.; Weinstein, I. B. Resveratrol induces growth inhibition, S-phase arrest, apoptosis, and changes in biomarker expression in several human cancer cell lines. *Clin. Cancer Res.* **2002**, *8*, 893-903.
168. Poussier, B.; Cordova, A.C.; Becquemin, J.P.; Sumpio, B.E. Resveratrol inhibits vascular smooth muscle cell proliferation and induces apoptosis. *J. Vasc. Surg.* **2005**, *42*, 1190-1197.
169. Larrosa, M.; Tomas-Barberan, F.A.; Espin, J.C. Grape polyphenol resveratrol and the related molecule 4-hydroxystilbene induce growth inhibition, apoptosis, S-phase arrest, and upregulation of cyclins A, E, and B1 in human SK-Mel-28 melanoma cells. *J. Agric. Food Chem.* **2003**, *51*, 4576-4584.
170. Meeran, S.M.; Katiyar, S.K. Grape seed proanthocyanidins promote apoptosis in human epidermoid carcinoma A431 cells through alterations in Cdk-cyclin cascade, and caspase-3 activation via loss of mitochondrial membrane potential. *Exp. Dermatol.* **2007**, *16*, 405-415.
171. Donnelly, L.E.; Newton, R.; Kennedy, G.E.; Fenwick, P.S.; Leung, R.H.; Ito, K.; Russell, R.E.; Barnes, P.J. Anti-inflammatory effects of resveratrol in lung epithelial cells: molecular mechanisms. *Am. J. Physiol. Lung Cell Mol. Physiol.* **2004**, *287*, L774–L783.
172. Rahman, I.; Kilty, I. Antioxidant therapeutic targets in COPD. *Curr. Drug Targets* **2006**, *7*, 707-720.
173. Xu, M.; Deng, B.; Chow, Y.L.; Zhao, Z.Z.; Hu, B. Effects of curcumin in treatment of experimental pulmonary fibrosis: a comparison with hydrocortisone. *J. Ethnopharmacol.* **2007**, *112*, 292-299.
174. Sakagami, Y.; Sawabe, A.; Komemushi, S.; All, Z.; Tanaka, T.; Iliya, I.; Inuma, M. Antibacterial activity of stilbene oligomers against vancomycin-resistant Enterococci (VRE) and methicillin-resistant Staphylococcus aureus (MRSA) and their synergism with antibiotics. *Biocontrol Sci.* **2007**, *12*, 7-14.
175. Furneri, P.M.; Piperno, A.; Sajia, A.; Bisignano, G. Antimycoplasmal activity of hydroxytyrosol. *Antimicrob. Agents Chemother.* **2004**, *48*, 4892-4894.
176. Nair, M.P.; Kandaswami, C.; Mahajan, S.; Nair, H.N.; Chawda, R.; Shanahan, T.; Schwartz, S.A. Grape seed extract proanthocyanidins downregulate HIV-1 entry coreceptors, CCR2b, CCR3 and CCR5 gene expression by normal peripheral blood mononuclear cells. *Biol. Res.* **2002**, *35*, 421-431.
177. Khanna, S.; Roy, S.; Bagchi, D.; Bagchi, M.; Sen, C.K. Upregulation of oxidant-induced VEGF expression in cultured keratinocytes by a grape seed proanthocyanidin extract. *Free Radic. Biol. Med.* **2001**, *31*, 38-42.
178. Ray, S.D.; Kumar, M.A.; Bagchi, D. A novel proanthocyanidin IH636 grape seed extract increases in vivo Bcl-XL expression and prevents acetaminophen-induced programmed and unprogrammed cell death in mouse liver. *Arch. Biochem. Biophys.* **1999**, *369*, 42-58.
179. Wong, M.C.; Portmann, B.; Sherwood, R.; Niemela, O.; Koivisto, H.; Parkkila, S.; Trick, K.; L'abbe, M.R.; Wilson, J.; Dash, P.R.; Srirajskanthan, R.; Preedy, V.R.; Wiseman, H. The cytoprotective effect of alpha-tocopherol and daidzein against d-galactosamine-induced oxidative damage in the rat liver. *Metabolism* **2007**, *56*, 865-875.

180. Kuzu, N.; Metin, K.; Dagli, A.F.; Akdemir, F.; Orhan, C.; Yalniz, M.; Ozercan, I.H.; Sahin, K.; Bahcecioglu, I.H. Protective role of genistein in acute liver damage induced by carbon tetrachloride. *Mediators Inflamm.* **2007**, *2007*, 36381.
181. Rucinska, A.; Kirko, S.; Gabryelak, T. Effect of the phytoestrogen, genistein-8-C-glucoside, on Chinese hamster ovary cells in vitro. *Cell Biol. Int.* **2007**, (in press).
182. Galati, G.; Sabzevari, O.; Wilson, J.X.; O'Brien, P.J. Prooxidant activity and cellular effects of the phenoxyl radicals of dietary flavonoids and other polyphenolics. *Toxicology* **2002**, *177*, 91-104.
183. Rietjens, I.M.C.M.; Boersma, M.G.; de Haan, L.; Spenkeliink, B.; Awad, H.M.; Cnubben, N.H.P.; van Zanden, J.J.; van der Woude, H.; Alink, G.M.; Koeman, J. H. The pro-oxidant chemistry of the natural antioxidants vitamin C, vitamin E, carotenoids and flavonoids. *Environ. Toxicol. Pharmacol.* **2002**, *11*, 321-333.
184. Huisman, A.; van de Wiel, A.; Rabelink, T.J.; van Faassen, E.E. Wine polyphenols and ethanol do not significantly scavenge superoxide nor affect endothelial nitric oxide production. *J. Nutr. Biochem.* **2004**, *15*, 426-432.
185. Sakihama, Y.; Cohen, M.F.; Grace, S.C.; Yamasaki, H. Plant phenolic antioxidant and prooxidant activities: phenolics-induced oxidative damage mediated by metals in plants. *Toxicology* **2002**, *177*, 67-80.
186. Fujisawa, S.; Atsumi, T.; Ishihara, M.; Kadoma, Y. Cytotoxicity, ROS-generation activity and radical-scavenging activity of curcumin and related compounds. *Anticancer Res.* **2004**, *24*, 563-569.
187. Nemeikaite-Ceniene, A.; Imbrasaitė, A.; Sergediene, E.; Cenas, N. Quantitative structure-activity relationships in prooxidant cytotoxicity of polyphenols: role of potential of phenoxyl radical/phenol redox couple. *Arch. Biochem. Biophys.* **2005**, *441*, 182-190.
188. Manach, C.; Scalbert, A.; Morand, C.; Rémésy, C.; Jiménez, L. Polyphenols: food sources and bioavailability. *Am. J. Clin. Nutr.* **2004**, *79*, 727-747.
189. Manach, C.; Williamson, G.; Morand, C.; Scalbert, A.; Rémésy, C. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am. J. Clin. Nutr.* **2005**, *81*, 230S-242S.
190. Gonthier, M.P.; Donovan, J.L.; Texier, O.; Felgines, C.; Rémésy, C.; Scalbert, A. Metabolism of dietary procyanidins in rats. *Free Radic. Biol. Med.* **2003**, *35*, 837-844.
191. Seeram, N.P.; Lee, R.; Heber, D. Bioavailability of ellagic acid in human plasma after consumption of ellagitannins from pomegranate (*Punica granatum L.*) juice. *Clin. Chim. Acta* **2004**, *348*, 63-68.
192. Chow, H.H.; Hakim, I.A.; Vining, D.R.; Crowell, J.A.; Ranger-Moore, J.; Chew, W.M.; Celaya, C.A.; Rodney, S.R.; Hara, Y.; Alberts, D.S. Effects of dosing condition on the oral bioavailability of green tea catechins after single-dose administration of Polyphenon E in healthy individuals. *Clin. Cancer Res.* **2005**, *11*, 4627-4633.
193. Henning, S.M.; Niu, Y.; Lee, N.H.; Thames, G.D.; Minutti, R.R.; Wang, H.; Go, V.L.; Heber, D. Bioavailability and antioxidant activity of tea flavanols after consumption of green tea, black tea, or a green tea extract supplement. *Am. J. Clin. Nutr.* **2004**, *80*, 1558-1564.

194. Henning, S.M.; Niu, Y.; Liu, Y.; Lee, N.H.; Hara, Y.; Thames, G.D.; Minutti, R.R.; Carpenter, C.L.; Wang, H.; Heber, D. Bioavailability and antioxidant effect of epigallocatechin gallate administered in purified form versus as green tea extract in healthy individuals. *J. Nutr. Biochem.* **2005**, *16*, 610-616.
195. Visioli, F.; Galli, C.; Bornet, F.; Mattei, A.; Patelli, R.; Galli, G.; Caruso, D. Olive oil phenolics are dose-dependently absorbed in humans. *FEBS Lett.* **2000**, *468*, 159-160.
196. Rasmussen, S.E.; Breinholt, V.M. Non-nutritive bioactive food constituents of plants: bioavailability of flavonoids. *Int. J. Vitam. Nutr. Res.* **2003**, *73*, 101-111.
197. Covas, M.I. Olive oil and the cardiovascular system. *Pharmacol. Res.* **2007**, *55*, 175-186.
198. Miro-Casas, E.; Covas, M.I.; Farre, M.; Fito, M.; Ortuño, J.; Weinbrenner, T.; Roset, P.; de la Torre, R. Hydroxytyrosol disposition in humans. *Clin. Chem.* **2003**, *49*, 945-952.
199. Caruso, D.; Visioli, F.; Patelli, R.; Galli, C.; Galli, G. Urinary excretion of olive oil phenols and their metabolites in humans. *Metabolism.* **2001**, *50*, 1426-1428.
200. Tuck, K.L.; Hayball, P.J.; Stupans, I. Structural characterization of the metabolites of hydroxytyrosol, the principal phenolic component in olive oil in rats. *J. Agric. Food Chem.* **2002**, *50*, 2404-2409.
201. Corona, G.; Tzounis, X.; Assunta Dessi, M.; Deiana, M.; Debnam, E.S.; Visioli, F.; Spencer, J.P. The fate of olive oil polyphenols in the gastrointestinal tract: implications of gastric and colonic microflora-dependent biotransformation. *Free Radic. Res.* **2006**, *40*, 647-658.
202. de la Torre-Carbot, K.; Jauregui, O.; Castellote, A.I.; Lamuela-Raventós, R.M.; Covas, M.I.; Casals, I.; López-Sabater, M.C. Rapid high-performance liquid chromatography-electrospray ionization tandem mass spectrometry method for qualitative and quantitative analysis of virgin olive oil metabolites in human low-density lipoproteins. *J. Chromatogr. A.* **2006**, *1116*, 69-75.
203. Carbonaro, M.; Grant, G.; Pusztai, A. Evaluation of polyphenol bioavailability in rat small intestine. *Eur. J. Nutr.* **2001**, *40*, 84-90.